

Crystal Structure of the High-temperature Solid Phases of Choline Tetrafluoroborate and Iodide

Hiroyuki Ishida^a, Hiroshi Ono^b, and Ryuichi Ikeda^b

^a Department of Chemistry, Faculty of Science, Okayama University, Okayama 700, Japan

^b Department of Chemistry, University of Tsukuba, Tsukuba 305, Japan

Z. Naturforsch. **52a**, 679–680 (1997);
received June 17, 1997

The crystal structure of the highest- and second highest-temperature solid phases of choline tetrafluoroborate and iodide was determined by X-ray powder diffraction. The structure in the highest-temperature phase of both salts is NaCl-type cubic ($a = 10.16(2) \text{ \AA}$, $Z = 4$ for tetrafluoroborate; $a = 10.08(2) \text{ \AA}$, $Z = 4$ for iodide). The second highest-temperature phase of tetrafluoroborate and iodide is CsCl-type cubic ($a = 6.198(6) \text{ \AA}$ and $Z = 1$) and tetragonal ($a = 8.706(2) \text{ \AA}$, $c = 6.144(6) \text{ \AA}$, and $Z = 2$), respectively. DSC was carried out for the iodide, where the presence of three solid-solid phase transitions was confirmed. Enthalpy and entropy changes of these transitions were evaluated.

Key words: Crystal structure; Powder X-ray diffraction; Phase transition.

In [1] we have shown that choline tetrafluoroborate, $[(\text{CH}_3)_3\text{NCH}_2\text{CH}_2\text{OH}]\text{BF}_4$, in the temperature range from the melting point at 485 K to 160 K has three solid phases named I, II, and III. In Phase I and II, stable between 485 and 402 K and between 402 and 268 K, respectively, isotropic rotation and translational self-diffusion of both cation and anion were observed by ^1H and ^{19}F NMR measurements. From the dynamical behaviour of the ions and the small entropy of fusion ($6.4 \text{ J K}^{-1} \text{ mol}^{-1}$) we concluded that this salt forms ionic plastic crystals in Phases I and II, the structures of which are expected to be of a high symmetry. A similar dynamical behaviour of the cation was observed in choline iodide, $[(\text{CH}_3)_3\text{NCH}_2\text{CH}_2\text{OH}]\text{I}$, by ^1H NMR [2–5]. Differential scanning calorimetry (DSC) [6] seemed to indicate that this salt undergoes two solid-solid phase transitions at 436 and 367 K with enthalpy changes of 6.23 and 12.8 kJ mol^{-1} , respectively. The isotropic rotation and self-diffusion of the cation in the highest-temperature phase [2, 3] and the second highest-temperature phase [3–5] (named Phase I and II, respec-

tively, in this text [7]) were detected by ^1H NMR. Furthermore, both salts show an unusual phenomenon, i.e. “quenching” of the cationic self-diffusion at the transition point from Phase II to Phase I [1, 3].

In the present study, X-ray powder diffractions in Phases I and II of the two salts were taken. In addition, DSC was carried out for the iodide to confirm the presence of a third highest-temperature phase existing between 366–360 K, as reported by Burnett et al. [5].

$[(\text{CH}_3)_3\text{NCH}_2\text{CH}_2\text{OH}]\text{BF}_4$ was prepared as described in [1]. $[(\text{CH}_3)_3\text{NCH}_2\text{CH}_2\text{OH}]\text{I}$ was purchased from Sigma Chemical Co. and recrystallized from ethanol. X-ray powder patterns were taken using a Philips X'pert PW3040/00 diffractometer. DSC was carried out using a Perkin-Elmer DSC7 in the range from 450 to 330 K.

The X-ray powder diffraction angles (2θ) in Phases I and II of tetrafluoroborate taken at ca. 420 and 300 K, respectively, are shown in Table 1. The structure of Phase I could be interpreted by an NaCl-type cubic lattice with $a = 10.16(2) \text{ \AA}$, $Z = 4$, $V = 1049(7) \text{ \AA}^3$ and $D_x = 1.209(8) \text{ Mg m}^{-3}$, and that of Phase II by a CsCl-type cubic lattice with

Table 1. Observed and calculated 2θ values of X-ray powder patterns in the highest- and second highest-temperature phases (Phase I and II) of $[(\text{CH}_3)_3\text{NCH}_2\text{CH}_2\text{OH}]\text{BF}_4$ taken at ca. 420 and 300 K, $\lambda(\text{Cu K}\alpha) = 1.5418 \text{ \AA}$, (Phase I: NaCl-type cubic, $a = 10.16(2) \text{ \AA}$, $Z = 4$, $V = 1049(7) \text{ \AA}^3$, and $D_x = 1.209(8) \text{ Mg m}^{-3}$; Phase II: CsCl-type cubic, $a = 6.198(6) \text{ \AA}$, $Z = 1$, $V = 238.1(7) \text{ \AA}^3$, $D_x = 1.332(4) \text{ Mg m}^{-3}$)

Observed		Calculated	
2θ (deg) (± 0.02)	I (%)	2θ (deg)	hkl
Phase I			
15.12	2	15.10	111
17.47	100	17.46	200
24.78	20	24.79	220
29.12	1	29.15	311
30.48	5	30.48	222
35.33	2	35.34	400
Phase II			
14.30	1	14.29	100
20.24	100	20.26	110
24.87	2	24.88	111
28.80	2	28.81	200
32.33	5	32.30	210

Reprint requests to Prof. Dr. H. Ishida; Fax: +81-86-251-8497.

0932-0784 / 97 / 0800-0679 \$ 06.00 © – Verlag der Zeitschrift für Naturforschung, D-72027 Tübingen



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

Table 2. Observed and calculated 2θ values of X-ray powder patterns in the highest- and second highest-temperature phases (Phase I and II) of $[(\text{CH}_3)_3\text{NCH}_2\text{CH}_2\text{OH}]\text{I}$ taken at ca. 460 and 400 K, λ ($\text{Cu K}\alpha$) = 1.5418 Å, (Phase I: NaCl-type cubic, $a = 10.08(2)$ Å, $Z = 4$, $V = 1024(7)$ Å³, and $D_x = 1.50(1)$ Mg m⁻³; Phase II: tetragonal, $a = 8.706(2)$, $c = 6.144(6)$ Å, $Z = 2$, $V = 465.7(7)$ Å³, $D_x = 1.648(3)$ Mg m⁻³)

Observed		Calculated	
2θ (deg)	I (%)	2θ (deg)	hkl
Phase I			
15.25	20	15.22	111
17.62	100	17.60	200
24.99	60	24.99	220
29.38	70	29.39	311
30.73	30	30.73	222
35.57	2	35.63	400
38.93	10	38.95	331
39.97	5	40.00	420
Phase II			
14.39	10	14.39	110
		14.42	001
17.69	7	17.67	101
20.42	100	20.40	200
		20.42	111
25.05	10	25.07	201
27.08	10	27.11	211
29.00	5	29.01	220
30.84	5	30.81	300
		30.87	102
32.49	10	32.52	310
35.72	7	35.74	311
42.81	1	42.83	410

$a = 6.198(6)$ Å, $Z = 1$, $V = 238.1(7)$ Å³ and $D_x = 1.332(4)$ Mg m⁻³. The "quenching" of the cationic self-diffusion in tetrafluoroborate, therefore, can be understood in terms of the change of lattice structure. The self-diffusion processes, in which the constituent ions in the CsCl-type structure migrate more easily than those in the NaCl-type, were discussed in [8, 9].

Table 2 shows the X-ray data in Phases I and II of the iodide obtained at ca. 460 and 400 K, respectively. The data of Phase I were assigned to an NaCl-type cubic lattice with $a = 10.08(2)$ Å,




Fig. 1. DSC curves recorded for $[(\text{CH}_3)_3\text{NCH}_2\text{CH}_2\text{OH}]\text{I}$ around 365 K with heating rates of 10 K min⁻¹ (a), 5 K min⁻¹ (b), 2 K min⁻¹ (c), and 1 K min⁻¹ (d).

$Z = 4$, $V = 1024(7)$ Å³ and $D_x = 1.50(1)$ Mg m⁻³ isomorphous with Phase I of the tetrafluoroborate, while that of Phase II was assigned to a tetragonal lattice with $a = 8.706(2)$, $c = 6.144(6)$ Å, $Z = 2$, $V = 465.7(7)$ Å³ and $D_x = 1.648(3)$ Mg m⁻³. The low symmetry of the structure in Phase II of the iodide is consistent with the fact that the activation energy of cationic self-diffusion in Phase II of the iodide is much larger than that of the tetrafluoroborate (105–110 kJ mol⁻¹ for the iodide [3, 5] and 62 kJ mol⁻¹ for the tetrafluoroborate [1]).

DSC measured on $[(\text{CH}_3)_3\text{NCH}_2\text{CH}_2\text{OH}]\text{I}$ with a heating rate of 2 K min⁻¹ showed three thermal anomalies attributable to solid-solid transitions at 362, 368, and 435 K, indicating clearly the existence of the fourth solid phase found by Burnett et al. [5]. Corresponding enthalpy (entropy) changes were evaluated to be 11.9 ± 0.1 (32.9 ± 0.3), 0.87 ± 0.04 (2.4 ± 0.1), and 6.19 ± 0.03 kJ mol⁻¹ (1.42 ± 0.07 J K⁻¹ mol⁻¹) in the same order. When the sample heating rate was raised, the anomaly peak at 362 K overlapped with the peak at 368 K, as shown in Figure 1. This may be the reason why the previous DSC showed only two solid-solid phase transitions.

- [1] H. Ishida, T. Asaji, Y. Furukawa, and R. Ikeda, Chem. Lett. 1207 (1993).
- [2] J. D. Graham and R. H. Hannon, J. Magn. Reson. **23**, 97 (1976).
- [3] C. A. McDowell, P. Raghunathan, and D. S. Williams, J. Chem. Phys. **66**, 3240 (1977).
- [4] J. D. Graham and R. H. Hannon, J. Chem. Phys. **64**, 1204 (1976).
- [5] L. J. Burnett, R. M. Knowles, and J. D. Graham, J. Chem. Phys. **68**, 2514 (1978).
- [6] V. Petrouleas and R. M. Lemmon, J. Chem. Phys. **69**, 1315 (1978).

- [7] Although Mc'Dowell et al. [3] designated the highest- and second highest-temperature solid phases of iodide as Phase III and II, respectively, we have labeled these phases as Phase I and II in the order of decreasing temperature.
- [8] J. F. Laurent and J. Bénard, J. Phys. Chem. Solids **3**, 7 (1957).
- [9] M. Tansho, Y. Furukawa, D. Nakamura, and R. Ikeda, Ber. Bunsenges. Phys. Chem. **96**, 550 (1992).