On the Observation of Decoherence with a Movable Mirror

R. Folman^{a,b}, J. Schmiedmayer^b, H. Ritsch^c, and D. Vitali^d

^a Institut f
ür Experimental Physik, Universit
ät Innsbruck, Technikerstra
ße 25, A-6020 Innsbruck, Austria

^b Physikalisches Institut, Universität Heidelberg, D-69120 Heidelberg, Germany

^c Institut für Theoretische Physik, Universität Innsbruck,

Technikerstraße 25, A-6020 Innsbruck, Austria

^d Dip. di Matematica e Fisica, and Unità INFM, Università di Camerino, I-62032 Camerino, Italy

Reprint requests to Dr. R. F.; E-mail: folman@physi.uni-heidelberg.de

Z. Naturforsch. 56 a, 140-144 (2001); received February 6, 2001

Presented at the 3rd Workshop on Mysteries, Puzzles and Paradoxes in Quantum Mechanics, Gargnano, Italy, September 17 - 23, 2000.

Recently it has been proposed to use parity as a measure of the mechanism behind decoherence or the transformation from quantum to classical. Here, we show that the proposed experiment is more feasible than previously thought, as even an initial thermal state would exhibit the hypothesized symmetry breaking. – Pacs: 03.75.Be, 03.65.Nk

Key words: Decoherence; Statistical Mixture; Parity Measurement; Symmetry Breaking; Movable Mirror.