

**Stereospecific
Four-Bond Phosphorus-Phosphorus Spin
Couplings in Phosphazeny-Phosphazenes**

M. BIDDLESTONE*, R. KEAT**, H. ROSE***,
D. S. RYCROFT**, and R. A. SHAW****

* Department of Chemistry, University of Khartoum,
P. O. Box 321, Khartoum, Sudan.

** Department of Chemistry, University of Glasgow,
Glasgow, G12 8 QQ, U. K.

*** Lehrstuhl für Anorganische Chemie I der Ruhr-
Universität Bochum, D-463 Bochum-Querenburg,
Postfach 2148, BRD.

**** Department of Chemistry, Birkbeck College,
(University of London), Malet Street, London
WC1E 7HX, U. K.


(Z. Naturforsch. **31b**, 1001-1002 [1976]; received March 4, 1976)

**³¹P NMR Spectra,
Four-Bond Phosphorus-Phosphorus Coupling,
Conformation, Phosphazenylicyclophosphazenes**

Four-bond phosphorus-phosphorus coupling constants have been measured from the ³¹P NMR spectra of phosphazenylicyclophosphazenes. Their magnitude appears to be related to the conformation adopted by the phosphazeny-yl-group relative to the phosphazene ring.

The two-bond spin-spin coupling constant, ²J_{PNP} is well documented¹ in both cyclic and acyclic phosphazenes. However, very few four-bond coupling constants, ⁴J_{P-N-P-N-P}, have been reported².

Examination of the ³¹P NMR spectra of a series of phosphazeny-yl-derivatives of cyclotriphosphazatrienes (**1**) revealed the presence of ⁴J_{P-N-P-N-P}.



The largest coupling (7.5 Hz) was in the trimethylphosphazeny-yl-derivative (**1a**)³ (for numbering see Table) and double and triple resonance experiments were carried out to establish its relative sign. The results are shown in the Figure, where a homonuclear ³¹P{³¹P} experiment (carried out under conditions of proton noise decoupling) showed that $J_{\underline{P}_A-\underline{N}-\underline{P}_C}$ and $J_{\underline{P}_A-\underline{N}-\underline{P}_C-\underline{N}-\underline{P}_D}$ have the same sign. In principle, the same experiment gives the relative

Requests for reprints should be sent to Dr. R. KEAT, Department of Chemistry, University of Glasgow, Glasgow, G12 8 QQ, U. K., or

Professor R. A. SHAW, Department of Chemistry, Birkbeck College, (University of London), Malet Street, London, WC1E 7HX, U. K.




Figure. ³¹P NMR spectra of **1a** (R = Me, X = Y = Cl), (a) normal spectrum with complete ¹H decoupling, (b) and (c) as in (a), but with irradiation of P_C signals (see text).

signs of $J_{\underline{P}_C-\underline{N}-\underline{P}_D}$ and $J_{\underline{P}_A-\underline{N}-\underline{P}_C-\underline{N}-\underline{P}_D}$, but the P_A resonance is broad (presumably because of quadrupolar relaxation of adjoining nitrogen and/or chlorine nuclei) and this combined with intensity changes, caused by population transfer effects, and the weakness of the available phosphorus irradiation field ($\frac{\gamma(^{31}\text{P})B_2}{2\pi} < 4 \text{ Hz}$) meant that interpretable effects could not be observed. However, selective irradiation of P_A and observation of P_D showed that $J_{\underline{P}_A-\underline{N}-\underline{P}_C}$ and $J_{\underline{P}_C-\underline{N}-\underline{P}_D}$ had the same sign. There is good reason to believe that $J_{\underline{P}-\underline{N}-\underline{P}}$ will be positive^{1,4}, so that the four bond coupling is also positive. This was confirmed by examination of the ¹H spectrum of this compound which consists of a doublet of doublets: ¹H{³¹P} double resonance showed that $J_{\underline{P}_C-\underline{N}-\underline{P}_D}$ and $J_{\underline{P}_C-\underline{N}-\underline{P}_D-\underline{C}-\underline{H}}$ have the same sign and that $J_{\underline{P}_C-\underline{N}-\underline{P}_D}$ and $J_{\underline{P}_D-\underline{C}-\underline{H}}$ are of opposite sign, the latter generally being found to be negative in phosphorus(V) compounds⁵. Homonuclear ³¹P{³¹P} experiments on the compounds^{3,6} listed in the Table showed that the four-bond couplings are positive, with the exception of that for **1e** where $J_{\underline{P}_A-\underline{N}-\underline{P}_C-\underline{N}-\underline{P}_D}$ is -0.4 Hz.

The hitherto reported P-N-P couplings in cyclophosphazenes are not simply related to the electro-negativity of the substituents on the phosphorus atoms¹ and the same generalisation may be made for $J_{\underline{P}_A-\underline{N}-\underline{P}_C}$ and $J_{\underline{P}_A-\underline{N}-\underline{P}_C-\underline{N}-\underline{P}_D}$ (Table). For example, the four-bond coupling becomes less positive in the order X = Cl > OEt > NMe₂ > Ph. However, there is evidence to suggest that this coupling constant is related to the preferred conformation adopted by the phosphazeny-yl-group relative to the phosphazene ring. The crystal structure of **1e**⁷ shows that the plane containing the exocyclic P-N-P unit lies perpendicular to the plane containing the phosphazene ring (type II conformation⁸), but in **1** (R = Ph, X = Cl, Y = Cl)⁹, and in 2-*trans*-6-N₄P₄Cl₆ (N = PPh₃)₂¹⁰, the Ph₃P=N-group is rotated by approximately 90° (type I

Table. ^{31}P NMR data^a for phosphazenylicyclophosphazenes.

Compound	R	X	Y	$\delta_{\text{P}}^{\text{b}}$	$J_{\text{P-N-P}}$ (Hz) ^c	$J_{\text{P-N-P-N-P}}$ (Hz) ^d	
1a	Me	Cl	Cl	A (= B)	20.9	$J_{\text{P}_A \dots \text{P}_C}$ 44.5	$J_{\text{P}_A \dots \text{P}_D} + 7.5^{\text{e}}$
				C	1.1	$J_{\text{P}_C \dots \text{P}_D}$ 16.0	
				D	25.4		
1b	Ph	Cl	Cl	A (= B)	20.3	$J_{\text{P}_A \dots \text{P}_C}$ 47.5	$J_{\text{P}_A \dots \text{P}_D} + 3.4$
				C	0.2	$J_{\text{P}_C \dots \text{P}_D}$ 27.8	
				D	15.4		
1c	Ph	OEt	Cl	A (= B)	18.5	$J_{\text{P}_A \dots \text{P}_C}$ 49.5	$J_{\text{P}_A \dots \text{P}_D} + 0.9$
				C	-1.6	$J_{\text{P}_C \dots \text{P}_D}$ 39.0	
				D	13.5		
1d	Ph	NMe ₂	Cl	A (= B)	17.1	$J_{\text{P}_A \dots \text{P}_C}$ 36.8	$ J_{\text{P}_A \dots \text{P}_D} < 0.1$
				C	4.3	$J_{\text{P}_C \dots \text{P}_D}$ 28.4	
				D	13.6		
1e	Ph	Ph	Cl	A (= B)	16.1	$J_{\text{P}_A \dots \text{P}_C}$ 21.9	$J_{\text{P}_A \dots \text{P}_D} - 0.4$
				C	2.9	$J_{\text{P}_C \dots \text{P}_D}$ 5.8	
				D	14.4		
1f	Ph	Cl	NMe ₂	A	21.4	$J_{\text{P}_A \dots \text{P}_B}$ 42.8	$J_{\text{P}_A \dots \text{P}_D} + 5.5$ $J_{\text{P}_B \dots \text{P}_D} + 3.3$
				B	26.4	$J_{\text{P}_A \dots \text{P}_C}$ 57.4	
				C	4.4	$J_{\text{P}_B \dots \text{P}_C}$ 57.0	
				D	14.6	$J_{\text{P}_C \dots \text{P}_D}$ 26.2	
2-trans-6-N ₄ P ₄ Cl ₆ (N=P _D Ph ₃) ₂				D	13.8		$ J_{\text{P}_D \text{-N-P-N-P} 4.6^{\text{f}}$

^a Obtained on a Varian XL-100 in the C. W. mode at 40.5 MHz in CDCl₃ solution at ca. 25°.

^b Relative to 85% H₃PO₄; low field shifts are positive.

^c All couplings are positive, ± 0.5 Hz.

^d ± 0.1 Hz.

^e $J_{\text{P}_D \text{-C-H}} = -13.02$ Hz, $J_{\text{P}_C \text{-N-P}_D \text{-C-H}} = +0.65$ Hz.

^f Apparent coupling.

conformation⁸; see ref.⁹). Furthermore, basicity measurements on a series of phosphazenylicyclophosphazenes indicate that electron supply from the Ph₃P=N-group to the cyclophosphazene ring is more effective in **1b** and **1f** than in **1d**, **1e** or **1c** (at least on protonation). The smaller, or least positive, four bond couplings may therefore be correlated with a preferred conformation in which the plane containing the exocyclic P-N-P unit is approximately perpendicular to the plane containing the phosphazene ring. The relatively inefficient electron supply to the cyclophosphazene ring in this conformation would be consistent with a reduced Fermi contact contribution to $J_{\text{P}_A \text{-N-P}_C \text{-N-P}_D}$. The

reduction in this coupling is paralleled by a progressive high-field shift for the P_A signals. This trend in ^{31}P shifts is often observed with increasing chlorine content of aminochlorocyclophosphazenes, where the phosphorus atoms become progressively more electron deficient.

It is interesting, but perhaps surprising, in view of the observations on $J_{\text{P}_A \text{-N-P}_C}$, to find that the positive two bond coupling, $J_{\text{P}_C \text{-N-P}_D}$, is linearly related to electronegativity parameters for the X-substituents in **1b-1e**.

R. K. and D. S. R. thank the S. R. C. (U. K.) for assistance in the purchase of NMR equipment.

¹ E. G. FINER and R. K. HARRIS, Progress in NMR Spectroscopy **6**, 86 [1970]; R. KEAT, R. A. SHAW, and M. WOODS, J. Chem. Soc. Dalton, in press.

² G. R. FEISTEL and T. MOELLER, J. Inorg. Nucl. Chem. **29**, 2731 [1967]; M. BIDDLESTONE and R. A. SHAW, J. Chem. Soc. (A) **1971**, 2715; H. W. ROESKY, Chem. Ber. **105**, 1439 [1972].

³ H. ROSE and R. A. SHAW, unpublished results.

⁴ P. CLARE, D. B. SOWERBY, R. K. HARRIS, and M. M. WAZEER, J. Chem. Soc. Dalton **1975**, 625.

⁵ W. MCFARLANE, Proc. Roy. Soc. (London) **A 306**, 185 [1968].

⁶ M. BIDDLESTONE and R. A. SHAW, J. Chem. Soc. Dalton **1973**, 2740.

⁷ M. BIDDLESTONE, G. J. BULLEN, P. E. DANN, and R. A. SHAW, Chem. Commun. **1974**, 56.

⁸ R. A. SHAW, Pure Appl. Chem. **44**, 317 [1975]; Z. Naturforsch. **31b**, 641 [1976].

⁹ Y. S. BABU, T. S. CAMERON, S. S. KRISHNAMURTHY, H. MANOHAR, and R. A. SHAW, Z. Naturforsch. **31b**, 999 [1976].

¹⁰ G. J. BULLEN and P. E. DANN, unpublished results.

¹¹ M. BIDDLESTONE, S. N. NABI, and R. A. SHAW, J. Chem. Soc. Dalton **1975**, 2634.