A New Prenylated Hydroquinone from the Roots of *Garcinia atroviridis* Griff ex T. Anders (Guttiferae)

Dharma Permana^a, Nordin Hj. Lajis^a, Khozirah Shaari^a, Abdul M. Ali^b, Mukram M. Mackeen^b, Mariko Kitajima^c, Hiromitsu Takayama^c, and Norio Aimi^c

- ^a Laboratory of Phytomedicines, Institute of Bioscience, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia
- ^b Department of Biotechnology, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia
- ^c Faculty of Pharmaceutical Sciences, Chiba University, I-33 Yayoi-cho, Inage-ku, Chiba 263-8522, Japan

Reprint requests to Prof. Dr. N. H. Lajis. E-mail: nhlajis@ibs.upm.edn.my

Z. Naturforsch. 58bb, 332–335 (2003); received October 21, 2002

A continuing study on the phytochemical constituents of *Garcinia atroviridis* Griff ex T. Anders (Guttiferae) has led to the isolation of a new prenylated hydroquinone from the roots of the plant. The compound has been elucidated to be 4-methylhydroatrovirinone, primarily by the use of gradient NMR and mass spectroscopy. The roots also yielded the known morelloflavone and its 7-O- β -D-glucopyranoside, fukugiside, together with 14-*cis*-do-cosenoic acid.

Key words: Prenylated Hydroquinone, Garcinia atroviridis