

# Isolation, Characterization and Crystal Structure of Natural Eremophilenolide from *Ligularia sagitta*

Lyi Li<sup>a</sup>, Li-Wen Xu<sup>b</sup>, Yan-Feng Jiang<sup>c</sup>, Chan-Juan Xi<sup>c</sup>, Han-Qing Wang<sup>a</sup>, and You-Rui Suo<sup>d</sup>

<sup>a</sup> Lanzhou Institute of Chemical Physics, Chinese Academy of Sciences, Lanzhou, 730000, China

<sup>b</sup> State key laboratory of Oxo Synthesis and Selective Oxidation, Lanzhou Institute of Chemical Physics, Chinese Academy of Sciences, Lanzhou, 730000, China

<sup>c</sup> Department of Chemistry, Tsinghua University, Beijing, 100084, China

<sup>d</sup> Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining, 810008, China

Reprint requests to Prof. H.-Q. Wang. E-mail: whqwt@hotmail.com. Fax: +86-931-8277088.

Z. Naturforsch. **59b**, 921 – 924 (2004); received January 8, 2004

A new eremophilenolide 1 $\beta$ ,10 $\beta$ -epoxy-6 $\beta$ -acetox-3 $\beta$ -angeloyloxy-8 $\beta$ -hydroxy-eremophil-7(11)-en-8,12 $\alpha$ -olide (**1**), together with liguhodgsonal (**2**), esculetin (**3**) and  $\beta$ -sitosterol (**4**), was isolated from the aerial parts of *Ligularia sagitta*. The structure of the new constituent (**1**) was elucidated by spectroscopic methods and confirmed by single-crystal X-ray diffraction.

*Key words:* *Ligularia sagitta*, Eremophilenolide, Single-Crystal, X-Ray Diffraction