

Two New Iridoid Glucosides from *Verbascum salviifolium* Boiss.

Zeliha S. Akdemir^a, I. Irem Tatli^a, Erdal Bedir^{b,c}, and Ikhlas A. Khan^b

^a Hacettepe University, Faculty of Pharmacy, Dept. of Pharmacognosy,
Sihhiye, Ankara 06100, Turkey

^b National Center For Natural Products Research Institute of Pharmaceutical Sciences,
University of Mississippi, University, Mississippi 38677, USA

^c Present Address: Ege University, Faculty of Engineering, Dept. of Bioengineering,
Bornova, Izmir 35100, Turkey

Reprint requests to Z. S. Akdemir. Fax: +90-312-3114777. E-mail: zakdemir@hacettepe.edu.tr

Z. Naturforsch. **60b**, 113 – 117 (2005); received September 25, 2003

From the aerial parts of the plant *Verbascum salviifolium*, two new iridoid glucosides, 6-*O*- β -D-glucopyranosylcatalpol (**1**) and 6-*O*-(6''-*O*-*trans*-*p*-hydroxycinnamoyl)- β -D-glucopyranosylaucubin (**2**) along with five known iridoid glycosides, 6-*O*- β -D-glucopyranosylaucubin (**3**), 6-*O*- α -L-rhamnopyranosylcatalpol (**4**), verbaspinoside [= 6-*O*-(2''-*O*-*trans*-cinnamoyl)- α -L-rhamnopyranosylcatalpol] (**5**), pulverulentoside I [= 6-*O*-(2''-*O*-*trans*-*p*-methoxycinnamoyl-3''-*O*-acetyl)- α -L-rhamnopyranosylcatalpol] (**6**), and buddlejoside A₈ [= 6-*O*-(4''-*O*-*trans*-3,4-dimethoxycinnamoyl)- α -L-rhamnopyranosylcatalpol] (**7**) were isolated. The structures of the new compounds were established on the basis of spectroscopic evidence.

Key words: *Verbascum*, Scrophulariaceae, Iridoid Glycosides, 6-*O*- β -D-Glucopyranosylcatalpol, 6-*O*-(6''-*O*-*trans*-*p*-Hydroxycinnamoyl)- β -D-glucopyranosylaucubin