

Some Chemical Constituents of *Terminalia glaucescens* and their Enzymes Inhibition Activity

Atta-ur-Rahman^a, Seema Zareen^a, M. Iqbal Choudhary^a, M. Nadeem Akhtar^a, Shahida Shujaat^a, and F. N. Ngounou^b

^a H. E. J. Research Institute of Chemistry, International Center for Chemical Sciences, University of Karachi, Karachi-75270, Pakistan

^b Department of Organic Chemistry, University of Yaounde 1, P. O. Box 812, Yaounde, Cameroon

Reprint requests to Prof. Dr. M. I. Choudhary. Fax: (92-21) 9243190-9243191.
E-mail: hej@cyber.net.pk

Z. Naturforsch. **60b**, 347 – 350 (2005); received September 9, 2004

A new triterpenoid, glaucinoic acid (2α , 3β , 19α , 24-tetrahydroxyolean-12-en-30-oic acid) (**1**) along with several known compounds, arjunic acid (**2**), arjungenin (**3**), sericoside (**4**), and friedelin (**5**) were isolated from the stem barks of *Terminalia glaucescens*. These compounds showed β -glucuronidase inhibitory activity. The structures were identified on the basis of spectroscopic techniques.

Key words: *Terminalia glaucescens*, Combretaceae, Glaucinoic Acid, β -Glucuronidase Inhibition