

3-Substituted Coumarins from the Twigs of *Rhadinothamnus rudis* ssp. *amblycarpus*

Corinne Girard^a, Delphine Roux^a, Frédéric Muyard^a, Michel Colombain^a, François Tillequin^b, Peter G. Waterman^c, and Françoise Bévalot^a

^a Laboratoire de Pharmacognosie, EA3921 Optimisation Métabolique et Cellulaire, UFR des Sciences Médicales et Pharmaceutiques, Place Saint-Jacques, 25030 Besançon Cedex, France

^b Laboratoire de Pharmacognosie de l'Université René Descartes, U.M.R./C.N.R.S. N° 8638, Faculté des Sciences Pharmaceutiques et Biologiques, 4 Avenue de l'Observatoire, 75006 Paris, France

^c Centre for Phytochemistry and Pharmacology, Southern Cross University, PO Box 157, Lismore, NSW 2480, Australia

Reprint requests to Dr. F. Muyard. Fax: 33 (0)3 81665568. E-mail: frederic.muyard@univ-fcomte.fr

Z. Naturforsch. **60b**, 561 – 564 (2005); received January 11, 2005

The twigs of *Rhadinothamnus rudis* ssp. *amblycarpus* (*Phebalium rude* ssp. *amblycarpum*) yielded three new 3-substituted coumarins: rudicoumarin A-C (**1**–**3**). Their structures were established on the basis of their NMR and mass spectral data. In addition, ten other known compounds were also isolated, including phebaclavin I, schinicoumarin, seselin, and seven linear furocoumarins.

Key words: *Rhadinothamnus rudis* ssp. *amblycarpus*, *Phebalium rude* ssp. *amblycarpum*, Rutaceae, Rudicoumarin A-C, Schinicoumarin