

Phenolic Glycosides from *Symplocos racemosa*

Viqar U. Ahmad, Muhammad Zubair, Muhammad A. Abbasi, Farzana Kousar,
Farman Ullah, Naheed Fatima, and Muhammad. I. Choudhary

H. E. J. Research Institute of Chemistry, International Center for Chemical Sciences, University of
Karachi, Karachi-75270, Pakistan

Reprint requests to Prof. V. U. Ahmad. E-mail: vuahmad@cyber.net.pk., zubairbr@yahoo.com

Z. Naturforsch. **60b**, 1101 – 1104 (2005); received May 22, 2005

The re-investigation of the chemical constituents of the bark of the stem of *Symplocos racemosa* Roxb. led to the isolation of two new phenolic glycosides, *Symconoside A* (**1**) and *Symconoside B* (**2**). The structures of the new compounds were determined by 1D and 2D-homonuclear and heteronuclear NMR spectroscopy, chemical evidences, and by comparison with the published data of the closely related compounds. The phenolic glycosides **1** and **2** displayed *in vitro* inhibitory activity against phosphodiesterase-I with the IC_{50} values of 158 ± 0.02 and $900 \pm 0.08 \mu\text{M}$, respectively.

Key words: *Symplocos racemosa*, Symplocaceae, Symconoside A, Symconoside B,
Phosphodiesterase I