## Iridoid Glucosides from Turkish Phlomis tuberosa

Ihsan Calis<sup>a</sup>, Hasan Kirmizibekmez<sup>a</sup>, Tayfun Ersoz<sup>a</sup>, Ali A. Dönmez<sup>b</sup>, Charlotte H. Gotfredsen<sup>c</sup>, and Søren R. Jensen<sup>c</sup>

- <sup>a</sup> Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University, TR-06100 Ankara, Turkey
- <sup>b</sup> Department of Biology, Faculty of Science, Hacettepe University, 06532 Ankara, Turkey

<sup>c</sup> Department of Chemistry, The Technical University of Denmark, Building 201, DK-2800 Lyngby, Denmark

Reprint requests to Dr. Hasan Kirmizibekmez. Fax: +90 312 311 4777. E-mail: hasankbekmez@yahoo.com

## Z. Naturforsch. 60b, 1295-1298 (2005); received July 22, 2005

From the aerial parts of *Phlomis tuberosa* a new iridoid glucoside, chlorotuberoside was isolated together with five known iridoid glucosides, lamalbide, shanzhiside methyl ester, 7-*epi*-phlomiol (= phloyoside I), sesamoside and 5-deoxysesamoside. Two known phenylethanoid glycosides, for-sythoside B, and decaffeoylacteoside, three known neolignan glycosides, dehydrodiconiferyl alcohol 9'-*O*- $\beta$ -D-glucopyranoside, dihydrodehydrodiconiferyl alcohol 9-*O*- $\beta$ -D-glucopyranoside and dihydrodehydrodiconiferyl alcohol 9'-*O*- $\beta$ -D-glucopyranoside, luteolin 7-*O*- $\beta$ -D-glucopyranoside, as well as  $\beta$ -sitosterol 3-*O*- $\beta$ -D-glucopyranoside and 1-methyl-*O*- $\beta$ -D-glucopyranoside were also obtained and characterized. The structures of the isolates were elucidated on the basis of spectroscopic data. The three known phenylethanoid glycosides, verbascoside, leucosceptoside A and martynoside were identified by TLC comparison with authentic compounds.

Key words: Phlomis tuberosa, Lamiaceae, Iridoid Glucosides, Chlorotuberoside