

Two New Glycosides from *Sanicula lamelligera*

Ling-Yun Zhou^a, Hai-Yang Liu^a, Bai-Bo Xie^a, Zhi-Hai Liu^b, and Chang-Xiang Chen^a

^a State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, The Chinese Academy of Sciences, Kunming, Yunnan 650204, P. R. China

^b Guiyang College of Traditional Chinese Medicine, Guiyang, Guizhou 550001, P. R. China

Reprint requests to Prof. C.-X. Chen. Fax: +86-871-5219934. E-mail: cxchen@mail.kib.ac.cn

Z. Naturforsch. **61b**, 607 – 610 (2006); received July 14, 2005

Two new glycosides, 21-*O*- β -D-glucopyranosyl-olean-12-ene-3 β , 16 α , 21 β , 22 α , 28-pentol-3-*O*- β -D-glucopyranosyl (1 \rightarrow 2)-[β -D-arabinopyranosyl (1 \rightarrow 3)]- β -D-glucopyranoside (**1**) and 4-*O*- β -D-glucopyranosyl rosmarinic acid (**3**), along with the known compound rosmarinic acid (**2**) were isolated from the whole plants of *Sanicula lamelligera* Hance. Their structures were elucidated by spectroscopic analysis and chemical means.

Key words: *Sanicula lamelligera*, Triterpenoid Saponin, 4-*O*- β -D-Glucopyranosyl, Rosmarinic Acid