

Two Prenylated Flavones from the Tree Bark of *Artocarpus lanceifolius*

Yana M. Syah^a, Sjamsul A. Achmad^a, Norio Aimi^b, Euis H. Hakim^a, Lia D. Juliawaty^a,
and Hiromitsu Takayama^b

^a Department of Chemistry, Institut Teknologi Bandung, Jalan Ganesha 10, Bandung 40132, Indonesia

^b Graduate School of Pharmaceutical Sciences, Chiba University, 1-33, Yayoi-cho, Inageku, Chiba 263-8522, Japan

Reprint requests to Prof. Dr. Euis H. Hakim. E-mail: euis@chem.itb.ac.id

Z. Naturforsch. **61b**, 1134 – 1137 (2006); received December 22, 2005

Two new prenylated flavones, artoindonesianins Z-1 (**1**) and Z-2 (**2**), together with two other known prenylated flavones, artobiloxantone (**3**) and cycloartobiloxanthone (**4**), had been isolated and identified from the chloroform extract of the tree bark of *Artocarpus lanceifolius*. The structures of these compounds were determined based on spectroscopic data, including UV, IR, 1-D and 2-D NMR, and mass spectra. The significance of the presence of the isolated compounds to the chemotaxonomy of *Artocarpus* is briefly discussed.

Key words: Artoindonesianins Z-1 and Z-2, *Artocarpus lanceifolius*, Chemotaxonomy