

Acylated Iridoid Glycosides from the Flowers of *Verbascum lasianthum* Boiss. ex Benth

I. Irem Tatli^a, Ikhlas A. Khan^b, and Zeliha S. Akdemir^c

^a Department of Pharmaceutical Botany, Faculty of Pharmacy, Hacettepe University, Sıhhiye, Ankara TR 06100, Turkey

^b National Center For Natural Products Research Institute of Pharmaceutical Sciences, University of Mississippi, University, Mississippi, 38677, USA

^c Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University, Sıhhiye, Ankara TR 06100, Turkey

Reprint requests to Dr. I. I. Tatli. E-mail: itatli@hacettepe.edu.tr

Z. Naturforsch. **61b**, 1183 – 1187 (2006); received December 2, 2005

Two new (**1**–**2**) and five known iridoid glycosides (**3**–**7**) as well as a known saponin (**8**) were isolated from the flowers of *Verbascum lasianthum* and their structures were determined as 6-*O*-(4'-*O*-*trans*-*p*-coumaroyl)- α -L-rhamnopyranosylaucubin (**1**), 6-*O*-(4'-*O*-*trans*-*p*-methoxycinnamoyl)- α -L-rhamnopyranosylaucubin (**2**), sinuatol (**3**), aucubin (**4**), geniposidic acid (**5**), catalpol (**6**), ajugol (**7**) and ilwensisaponin A (**8**) on the basis of 1D and 2D NMR spectral analysis.

Key words: *Verbascum lasianthum*, Scrophulariaceae, Acylated Iridoid Glycosides, Lasianthoside I, Lasianthoside II