

# Two New Oleanane Triterpene Glycosides from the Bark of *Terminalia arjuna*

Asif Ali<sup>a</sup>, Mohammed Ali<sup>b</sup>, and Mohd S. Alam<sup>c</sup>

<sup>a</sup> Bioorganic Chemistry Laboratory, National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi-110067, India

<sup>b</sup> Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Hamdard University, New Delhi-110062, India

<sup>c</sup> Department of Chemistry, Faculty of Science, Hamdard University, New Delhi-110062, India

Reprint requests to Prof. Dr. M. Ali. E-mail: mali\_chem@yahoo.co.in

Z. Naturforsch. **61b**, 1282 – 1286 (2006); received February 28, 2005

Two new oleanane triterpene glycosides designated as termiarjunoside I (**1**) and termiarjunoside II (**2**), isolated from the bark of *Terminalia arjuna* (Combretaceae), have been characterized as olean-1 $\alpha$ ,3 $\beta$ ,9 $\alpha$ ,22 $\alpha$ -tetraol-12-en-28-oic acid-3 $\beta$ -D-glucopyranoside (**1**) and olean-3 $\alpha$ ,5 $\alpha$ ,25-triol-12-en-23,28-dioic acid-3 $\beta$ -D-glucopyranoside (**2**), respectively, on the basis of chemical and spectral data evidences.

*Key words:* *Terminalia arjuna*, Combretaceae, Termiarjunoside I, Termiarjunoside II