Synthesis and Crystal Structure of Schiff Bases Based on AMTTO (AMTTO = 4-Amino-6-methyl-3-thio-3,4-dihydro-1,2,4-triazin-5(2*H*)-one)

Maasoomeh Tabatabaee<sup>a</sup>, Mitra Ghassemzadeh<sup>b</sup>, Behnaz Zarabi<sup>a</sup>, and Bernhard Neumiiller<sup>c</sup>

Reprint requests to Dr. M. Tabatabaee. E-mail: tabatabaee45m@yahoo.com

Z. Naturforsch. **61b**, 1421 – 1425 (2006); received May 23, 2006

Reaction of 4-amino-6-methyl-3-thio-3,4-dihydro-1,2,4-triazin-5(2*H*)-one (1) with 2-methoxy-benzaldehyde, 4-methoxybenzaldehyde, 2-pyridinecarbaldehyde, and 2,4-dichlorobenzaldehyde under classical heating and microwave irradiation in a solventfree system led to the corresponding imines (Schiff bases) 2–5. All synthesized compounds have been characterized by IR and NMR spectroscopy, mass spectrometry and by X-ray diffraction studies.

Key words: 1,2,4-Triazine, 1,2,4-Triazole, Schiff Base, Microwave Irradiation

<sup>&</sup>lt;sup>a</sup> Department of Chemistry, Islamic Azad University, Yazd Branch, Tehran, Iran

<sup>&</sup>lt;sup>b</sup> Chemistry and Chemical Engineering Research Center of Iran, P.O. Box 14335-186, Tehran, Iran <sup>c</sup> Fachbereich Chemie der Universität Marburg, Hans-Meerwein-Straße, 35032 Marburg, Germany