

Alkaloids from *Palicourea coriacea* (Cham.) K. Schum.

Claudia A. do Nascimento^a, Mislene S. Gomes^a, Luciano M. Lião^a,
Cecília M. A. de Oliveira^a, Lucilia Kato^a, Cleuza C. da Silva^b, and Clara M. A. Tanaka^b

^a Instituto de Química, Universidade Federal de Goiás, Campus Samambaia, C.P. 131,
74001-970. Goiânia, GO – Brazil

^b Departamento de Química, Universidade Estadual de Maringá, Av. Colombo, 5790,
87020-900. Maringá, PR – Brazil

Reprint requests to Prof. Dr. Luciano M. Lião. E-mail: luciano@quimica.ufg.br.
Fax: +55 62 35211167

Z. Naturforsch. **61b**, 1443 – 1446 (2006); received January 30, 2006

Two new β -carboline alkaloids, *epi*-strictosidinic acid (**1**) and strictosidinic ketone (**2**), together with two known alkaloids, calycanthine (**3**) and strictosidinic acid (**4**) besides the triterpene ursolic acid (**5**) were isolated from the roots and leaves of *Palicourea coriacea*. The structures of the isolates were elucidated on the basis of spectral data, particularly COSY, HMQC and HMBC experiments, and HR mass spectra.

Key words: Alkaloid, β -Carbolines, *Palicourea coriacea*, *epi*-Strictosidinic Acid,
Strictosidinic Ketone