

Butyrospermol Fatty Acid Esters from the Fruit of a Chinese Mangrove *Xylocarpus granatum*

Jun Wu^a, Minyi Li^a, Zhihui Xiao^a, and Yuan Zhou^b

^a Guangdong Key Laboratory of Marine Materia Medica, South China Sea Institute of Oceanology, Chinese Academy of Sciences, 164 West Xingang Road, Guangzhou 510301, P. R. China

^b Chemistry & Life Science College, China Three Gorges University, 8 University Road, Yichang 443002, P. R. China

Reprint requests to Dr. Jun Wu. Fax: +86-20-84451672. E-mail: wwujun2003@yahoo.com

Z. Naturforsch. **61b**, 1447 – 1449 (2006); received April 24, 2006

A mixture of four new butyrospermol and two known β -sitosterol fatty acid esters, was obtained from the fruit of a Chinese mangrove *Xylocarpus granatum*. The structures of these compounds were established on the basis of spectroscopic data and chemical means. The butyrospermol fatty acid esters were characterized as mainly comprising butyrospermol 3β -*O*-palmitate (**1**), butyrospermol 3β -*O*-oleate (**2**), butyrospermol 3β -*O*-stearate (**3**) and butyrospermol 3β -*O*-linoleate (**4**). The β -sitosterol fatty acid esters were identified as mainly containing β -sitosterol 3β -*O*-myristate and β -sitosterol 3β -*O*-oleate.

Key words: Butyrospermol Fatty Acid Ester, *Xylocarpus granatum*