

Xylodiol, a New Atisane Diterpenoid from *Xylophia Langsdorffiana* St.-Hil. & Tul. (Annonaceae)

Josean Fechine Tavares^a, Marianna Vieira B. Silva^a, Karine F. Queiroga^a,
Margareth F. F. Melo Diniz^a, José M. Barbosa Filho^a, Marcela Haun^b,
Patrícia S. Melo^{b,c}, and Marcelo Sobral da Silva^a

^a Laboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, CP 5009, João Pessoa,
PB 58051-970, Brazil

^b Departamento de Bioquímica, Instituto de Biologia, Universidade Estadual de Campinas, CP 6109,
Campinas, SP 13083-970, Brazil

^c Faculdades Integradas Metropolitanas de Campinas, Rua Abolição, nº 1827, Swift, Campinas, SP,
Brazil

Reprint requests to Dr. Marcelo Sobral da Silva. E-mail: marcelosobral@lrf.ufpb.br

Z. Naturforsch. **2007**, *62b*, 742–744; received November 20, 2006

Two diterpenes were isolated from the leaves of *Xylophia langsdorffiana*, the new *ent*-atisane-7 α ,16 α -diol named xylodiol (**1**) and the known 8 (17),12E,14-labdatrien-18-oic acid (**2**). The structures of these isolates were deduced by spectroscopic data interpretation. Compound **2** was less toxic than xylodiol on V79 cells and hepatocytes.

Key words: *Xylophia langsdorffiana*, Annonaceae, Atisane, Labdane, Cytotoxicity