

Two New Iridoid Glycosides from *Lagotis yunnanensis*

Xiao-Dong Yang^a, Li-Juan Yang^b, Shu Yang^a, Jing-Feng Zhao^a, Hong-Bin Zhang^a, and Liang Li^a

^a Key Laboratory of Medicinal Chemistry for Natural Resources (School of Chemical Science and Technology, Yunnan University), Ministry of Education, Kunming 650091, P.R. China

^b School of Chemistry and Bio-Science, Yunnan Nationalities University, Kunming 650030, P. R. China

Reprint requests to Prof. Liang Li. Fax: +86-871-5035538. E-mail: xdyang120@hotmail.com

Z. Naturforsch. **2007**, 62b, 749–752; received November 14, 2006

Two new iridoid glycosides, named lagotisoside D (**1**) and E (**2**), were isolated from *Logatis yunnanensis*. The structures of the two compounds were established as 6-*O*- α -L-(4''-*O*-*E*-3''',4'''-dimethoxycinnamoyl)rhamnopyranosylcatalpol (**1**) and 6-*O*- α -L-(4''-*O*-*Z*-3''',4'''-dimethoxycinnamoyl)rhamnopyranosylcatalpol (**2**) on the basis of spectral analysis.

Key words: *Logatis yunnanensis*, Iridoid Glycosides, Lagotisoside D, Lagotisoside E