

Chemical Constituents from the Roots and Rhizomes of *Clematis hexapetala* Pall.

Cai-xia Dong^{a,b}, She-po Shi^a, Ke-si Wu^b, and Peng-fei Tu^a

^a School of Pharmaceutical Sciences, Peking University Health Science Center, Beijing 100083, China

^b Shaanxi University of Chinese Medicine, Shaanxi, Xianyang, 712046, China

Reprint requests to Prof. Peng-fei Tu. Tel/Fax: +86-10-82802750. E-mail: pengfeitu@bjmu.edu.cn

Z. Naturforsch. **2007**, 62b, 854–858; received November 1, 2006

A new phenolic glycoside, 2,6-dimethoxy-4-(3-hydroxy-propen-1-yl)phenyl-4-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (**1**) and a new D-ribono- γ -lactone derivative, 3-*O*- β -D-glucopyranosyl-2-hydroxymethyl-D-ribono- γ -lactone (**3**), together with thirteen known compounds have been isolated from the roots and rhizomes of *Clematis hexapetala* Pall. The characterization of these compounds was achieved by various chromatographic and spectroscopic methods.

Key words: *Clematis hexapetala*, Chemical Constituents, Structure Elucidation