Diabietane Ether, a New Dimeric Abietane with an Ether Linkage from *Taxus cuspidata* Needles

Zhi-Yu Ni[a], Yi-Bing Wu[b], Mei Dong[a], Man-Li Zhang[b], Yu-Fang Wang[b], Françoise Sauriol[c], Chang-Hong Huo[b], Qing-Wen Shi[b,d], Yu-Cheng Gu[b,e], Hiromasa Kiyota[f], and Bin Cong[a,d]

[a] College of Basic Medicine, Hebei Medical University, Shijiazhuang, 050017, Hebei Province, P. R. China
[b] School of Pharmaceutical Sciences, Hebei Medical University, Shijiazhuang, 050017, Hebei Province, P. R. China
[c] Department of Chemistry, Queen's University, Kingston, Ontario, K7L 3N6, Canada
[d] Hebei Key Laboratory of Forensic Medicine, Shijiazhuang, 050017, Hebei Province, P. R. China
[e] Syngenta Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY, UK
[f] Graduate School of Agricultural Science, Tohoku University, Aoba-ku, Sendai 981-8555, Japan

Reprint requests to Q.-W. Shi.
E-mail: shiqingwen@hebmu.edu.cn or
H. Kiyota. E-mail: kiyota@biochem.tohoku.ac.jp

Z. Naturforsch. **2011**, *66b,* 1083 – 1086;
received July 20, 2011

Diabietane ether (**1**), a new abietane dimer connected by an ether linkage, and two known abietane diterpenoids (**2** and **3**) were isolated from the needles of *Taxus cuspidata.*

Key words: *Taxus cuspidata*, Abietanes, Diterpenes, Dimer, Structure Elucidation