Identification of Cyclic Enolethers from Insects: Alkyldihydropyranes from Bees and Alkyldihydro-4H-pyran-4-ones from a Male Moth*

W. Francke **, W. Mackenroth, W. Schröder, and S. Schulz

Institut für Organische Chemie der Universität, Martin-Luther-King-Platz 6, D-2000 Hamburg 13

J. Tengö

Ecological Station, Uppsala University, S-38600 Färjestaden

E. Engels and W. Engels ***

Institut für Biologie III (Zoologie) der Universität, Auf der Morgenstelle 28, D-7400 Tübingen

R. Kittmann, D. Schneider

Max-Planck-Institut für Verhaltensphysiologie, D-8131 Seewiesen

Z. Naturforsch. **40 c**, 145–147 (1985); received December 8, 1984

Mass spectrometric fragmentation patterns of alkyl-3,4dihydro-2H-pyranes and alkyl-2,3-dihydro-4H-pyran-4ones are described. Through GC/MS analyses, respective compounds showing unbranched carbon skeletons are identified for the first time as volatile signals of social and solitary bees and of the male moth *Hepialus hecta* L.

We studied mass spectrometric fragmentation patterns of cyclic enolethers to facilitate their identification from natural material [1]. Mass spectra of alkyl-3,4-dihydro-2 H-pyranes (1) are characterized by fragments which correspond to:

- A) α -cleavage at 2-C;
- B) retro cleavage ("Retro-Diels-Alder-Reaction");
- C) retro cleavage with hydrogen transfer to the oxygen containing fragment;
- D) an acylium fragment which contains 6-C and the respective substituent.

Signals caused by fragmentations C and D are particularly intense when the substituent at 6-C is a methyl- or an ethyl-group, while A and B are of low intensities when 6-C carries a longer chain.

* Dedicated to Prof. H. Francke-Grosmann on the occasion of her 85th birthday.

** Reprint requests to Dr. W. Francke.

*** In cooperation with the Departments of Genetics and Biology, University of São Paulo at Ribeirão Preto/Brazil. 0341-0382/85/0100-0145 \$ 01.30/0 We now found, that an alkyl chain of at least *n*propyl, attached to the sp²-carbon atom which carries the oxygen, in dihydropyranes as well as in dihydrofuranes and tetrahydrooxepines may furnish intense signals corresponding to:

- E) Mc-Lafferty rearrangement at the side chain;
- F) a "formal" γ -cleavage at the side chain;
- G) Mc-Lafferty rearrangement of the oxygen containing fragment produced by *retro* cleavage B.



Mass spectra of alkyl-2,3-dihydro-4H-pyran-4ones (2) resemble those of alkyl-3,4-dihydro-2Hpyranes because they also show fragmentations A-G. However, due to the presence of the carbonyl group dihydropyrones may give some additional signals.

On the basis of these results we identified for the first time several alkyldihydropyranes and alkyldihydro-4 H-pyran-4-ones both from social and solitary bees and from a male moth (see Table I).

During our investigations on odour communication in bees we studied the cephalic secretions of solitary bees, of social stingless bees and of honey bees. Among the complex multicomponent mixtures we identified small amounts of several new cyclic enolethers. In the abdomina of workers of Apis mellifica L. we found 2,6-dimethyl-3,4-dihydro-2Hpyrane (1a). This compound is also present in head extracts of the stingless bee Scaptotrigona bipunctata (Lepeletier) which additionally contain two bishomologues, 2-methyl-6-propyl-3,4-dihydro-2H-pyrane (1b) and 2-methyl-6-pentyl-3,4-dihydro-2H-pyrane (1 c). The latter compound was also found in Nannotrigona testaceicornis (Lepeletier), Plebeia droryana Friese, Tetragona clavipes (F.) and in some Taeniandrena species. Fig. 1 shows a plotted mass spectrum of 1c which we now identified as one of the main components of the cephalic scent mark secretion of the solitary bee Andrena wilkella K. [2]. Besides 6-butyl-3,4-dihydro-2H-pyrane (1f), head extracts of Partamona cupira (Smith) contain 6-heptyl-2methyl-3,4-dihydro-2H-pyrane (1d) and 2-methyl-6nonyl-3,4-dihydro-2H-pyrane (1e).

Compounds 1a-1e form a new row of unbranched bishomologue cryptic 2-hydroxyalkan-6-ones which seem to originate from the acetate pool and which



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz. This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung "Keine Bearbeitung") beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen. On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

No.	R ₁ CH ₃	R ₂ CH ₃	Insect species A. mellifica ^a S. bipunctata	MS (EI, 70 eV) m/z (%), [Fragmentation] 112 (20), 97 (10) [A], 83 (3), 71 (40) [C], 69 (12), 67 (6), 58 (8), 55 (31), 43 (100) [D]	¹ H-NMR 270 MHz	
1a					C ₆ D ₆ :	1.14 (d, 3 H), 1.41 (m, 2 H), 1.75 (s, 3 H), 1.85 (m, 2 H), 3.75 (m, 1 H), 4.43 (m, 1 H)
1b	CH ₃	C ₃ H ₇	S. bipunctata	140 (20), 125 (7) [A], [F], 112 (41) [E], 99 (18) [C], 97 (29), 83 (20), 71 (46) [D], 70 (22) [G], 55 (100), 43 (67)	C ₆ D ₆ :	0.91 (t, 3 H), 1.16 (d, 3 H), 1.26–1.52 (m, 2 H), 1.62 (m, 2 H), 1.77–2.04 (m, 2 H), 2.09 (t, 2 H), 3.78 (m, 1 H), 4.48 (m, 1 H)
1c	CH ₃	C ₅ H ₉	S. bipunctata N. testaceicornis P. droryana T. clavipes Taeniandrena spp.	168 (18), 126 (14) [B], 125 (41) [F], 112 (100) [E], 97 (36), 84 (29), 83 (46), 70 (35) [G], 55 (86), 43 (56)	C ₆ D ₆ :	0.88 (t, 3 H), 1.17 (d, 3 H), 1.23–1.67 (m, 8 H), 1.79–2.07 (m, 2 H), 2.13 (t, 2 H), 3.89 (m, 1 H), 4.15 (m, 1 H)
1d	CH3	C ₇ H ₁₅	P. cupira	196 (10), 125 (53) [F], 112 (100) [E], 97 (22), 84 (21), 83 (32), 70 (20) [G], 58 (18), 55 (60), 43 (37)	C ₆ D ₆ :	0.88 (t, 3 H), 1.18 (d, 3 H), 1.20-1.34 (m, 10 H), 1.55-1.71 (m, 2 H), 1.78-2.10 (m, 2 H), 2.16 (t, 2 H), 3.81 (m, 1 H), 4.53 (m, 1 H)
1e	CH ₃	C ₉ H ₁₇	P. cupira	224 (6), 125 (46) [F], 112 (100) [E], 97 (15), 84 (13), 83 (20), 70 (13) [G], 58 (13), 55 (38), 43 (27)	C ₆ D ₆ :	0.91 (t, 3 H), 1.18 (d, 3 H), 1.20-1.53 (m, 16 H), 1.78-2.06 (m, 2 H), 2.16 (t, 2 H), 3.79 (m, 1 H), 4.52 (m, 1 H)
lf	Н	C ₄ H ₉	P. cupira	140 (17), 111 (9) [F], 98 (100) [E], 85 (11) [D], 83 (27), 70 (20) [G], 57 (15), 56 (15), 55 (61), 43 (48)	C ₆ D ₆ :	0.88 (t, 3 H), 1.23–1.48 (m, 4 H), 1.78 (m, 2 H), 1.96 (m, 4 H), 3.96 (m, 2 H), 4.44 (t, 1 H)
2a	CH ₃	C_2H_5	H. hecta	140 (55), 125 (2) [A], 111 (3), 99 (92) [C], 98 (14) [B], 69 (100), 57 (62) [D], 55 (7), 43 (25)	CDCl ₃ :	1.08 (t, 3 H), 1.41 (d, 3 H), 2.22 (q, 2 H), 2.34 (m, 2 H), 4.44 (m, 1 H), 5.25 (m, 1 H)
2b	C_2H_5	C ₂ H ₅	H. hecta	154 (16), 126 (3), 125 (4) [A], 99 (100) [C], 98 (5) [B], 69 (51), 57 (51) [D], 56 (17), 43 (38)	CDCl ₃ :	0.99 (t, 3 H), 1.10 (t, 3 H), 1.62 – 1.89 (m, 2 H), 2.25 (q, 2 H), 2.37 (m, 2 H), 4.26 (m, 1 H), 5.28 (m, 1 H)
2c	C_2H_5	CH3		140 (35), 125 (3), 112 (3), 111 (3) [A], 97 (2), 85 (100) [C], 84 (4) [B], 69 (23), 56 (13), 43 (77) [D]	CDCl ₃ :	0.92 (t, 3 H), 1.55–1.83 (m, 2 H), 1.91 (s, 3 H), 2.28 (m, 2 H), 4.19 (m, 1 H), 5.18 (m, 1 H)
2d	CH ₃	C ₅ H ₉		182 (27), 167 (1) [A], 141 (58) [C], 140 (16) [B], 139 (17) [F], 126 (60) [E], 99 (11) [D], 98 (21), 97 (22), 84 (100) [G], 69 (72), 55 (25), 43 (31)	CDCl ₃ :	0.92 (t, 3 H), 1.34 (m, 4 H), 1.46 (d, 3 H), 1.58 (m, 2 H), 2.24 (t, 2 H), 2.42 (m, 2 H), 4.50 (m, 1 H), 5.31 (m, 1 H)

Table I. Cyclic enolethers identified from insects; mass spectra; ¹H-NMR-spectra.

^a Found in abdomina: has been erroneously reported as a constituent of the cephalic secretion¹.

Notizen

146







Fig. 2. 70 eV Mass spectrum of 6-ethyl-2-methyl-2,3-dihydro-4H-pyran-4-one.

may be biogenetically closely related to spiroacetals like 2,8-dimethyl-6,8-dioxaspiro [5.5]undecane which we also found in several of the insect species mentioned above [2, 3].

2,3-Dihydro-4H-pyranes are very rare in insects: 2,5-diethyl-2-formyl-2,3-dihydro-4H-pyrane, the Diels-Alder-dimer of 2-methylenebutanal, has been described from Blattidae [4].

As part of our studies on pheromones of male lepidoptera we analyzed the fruitlike smelling volatiles of the odoriferous organs of the male swift moth *Hepialus hecta* L. The gaschromatogram showed a major component and two minor ones, while several additional substances could be detected in trace amounts. The main compound (mol. weight $140 = C_8H_{12}O_2$ as determined by high resolution mass spectrometry) was identified as 6-ethyl-2-methyl-2,3-dihydro-4H-pyran-4-one (**2a**). Fig. 2

- W. Francke and W. Mackenroth, Angew. chem. Int. Ed. 21, 698 (1982).
- [2] G. Bergström, J. Tengö, W. Reith, and W. Francke, Z. Naturforsch. 37 c, 1124 (1982).
- [3] W. Francke, Nova Acta Reg. Soc. Scient. Upsaliensis Ser. V: C3, 119 (1984).
- [4] B. E. Wallbank and D. F. Waterhouse, J. Insect Physiol. 16, 2081 (1970).
- [5] V. Sinnwell, S. Schulz, W. Francke, R. Kittmann, and D. Schneider, Tetrahedron Letters 25 (1985) in press.

shows a plotted mass spectrum of this compound, which to our knowledge is the first compound identified from Hepialidae. Additionally, a homologue of (2a), 2,6-diethyl-2,3-dihydro-4H-pyran-4one (2b), proved to be present in trace amounts. The two minor components and some of the trace components mentioned above, are derivatives of the 2,9-dioxabicyclo [3.3.1]non-7-ene system [5].

While the new 2,3-dihydro-4H-pyran-4-ones show unbranched carbon skeletons, tetrasubstituted compounds which are probably derived from propionate units have been identified from *Stegobium paniceum* L. and *Lasioderma serricorne* F. (Col. Anobiidae) [6, 7].

GC analyses and GC/MS investigations were carried out on 50 m glas capillary columns with WG 11 as a stationary phase and on 50 m fused silica capillaries coated with SE 54. Mass spectra were obtained with a Varian MAT 311A. Chemical structures of natural products were confirmed by comparison of GC/MS data with those of authentic reference samples. Mass spectral data and ¹H-NMR data of the compounds are compiled in Table I.

Alkyl-3,4-dihydro-2 H-pyranes were prepared by Grignard reaction of lactones with alkylmagnesiumhalides followed by elimination of water from the obtained cyclic hemiacetals [8]. Optically active dihydropyranes may be produced from respective optically active lactones. Alkyl-2,3-dihydro-4 Hpyran-4-ones were prepared by acylation of β -ketoesters with $\alpha\beta$ -unsaturated acylhalogenides followed by cyclization, saponification and decarboxylation [9]. Bioassays with synthetic compounds will be described elsewhere.

Acknowledgements

The authors greatfully acknowledge financial support by the Deutsche Forschungsgemeinschaft, Fonds der Chemischen Industrie and the Swedish National Science Foundation.

- [6] Y. Kuwahara, H. Fukami, R. Howard, S. Ishii, F. Matsumura, and W. E. Burkholder, Tetrahedron 34, 1769 (1978).
- [7] T. Chuman, K. Mochizuki, K. Kato, M. Ono, and A. Okubo, Agric. Biol. Chem. 47, 1413 (1983).
- [8] D. Taub, N. N. Gerotra, R. D. Hofsommer, C. H. Kuo, H. L. Slates, S. Weber, and N. L. Wendler, Tetrahedron 24, 2443 (1968).
- [9] S. Gelin and R. Gelin, Bull. Soc. Chim. France 1968, 288.