

(Z)-3-Tetradecenyl Acetate as a Sex-Attractant Component in Gelechiinae and Anomologinae (Lepidoptera: Gelechiidae)

Ernst Priesner

Max-Planck-Institut für Verhaltensphysiologie,
D-8131 Seewiesen

Z. Naturforsch. **42c**, 1352–1355 (1987);
received August 25, 1987

Sex-Attractant, Attraction-Inhibitor, Δ 3-Alkenyl
Acetates, *Chionodes*, *Monochroa*, *Argolamprotes*,
Aproaerema, Gelechiidae

The title compound, unreported as an insect pheromone component, effectively attracted certain male Gelechiidae (genera *Chionodes*, *Monochroa*, *Argolamprotes*) as a single chemical. Trap captures with this chemical decreased on addition of either (*E*)-3-dodecenyl acetate, (*E*)-3-tetradecenyl acetate or (*Z*)-3-tetradecen-1-ol, the sexual attractants of other, closely related species. Results on an *Aproaerema* test species showing a synergistic attraction response to combinations of (*Z*)-3-tetradecenyl acetate with its homologue (*Z*)-3-dodecenyl acetate are included.

Long-chain alkenyl acetates are the most common structural type among identified sex pheromone and attractant components in the Lepidoptera [1]. The title compound, (*Z*)-3-tetradecenyl acetate = Z3-14:Ac, recently joined this group of semiochemicals when Willemse *et al.* [2] reported its attractiveness towards male *Chionodes electella* (Zell.) and *Monochroa tenebrella* (Hb.), members of the Gelechiidae subfamilies Gelechiinae and Anomologinae. Their report prompts a brief presentation of my own results obtained on this same and some structurally related compounds. The results define more closely the sex-attractant systems for the two species, demonstrate the attractiveness of Z3-14:Ac for further spp. of the same subfamilies, and provide evidence for involvement of this compound in synergistic attractant systems.

The response of certain male Gelechiidae to sources of Z3-14:Ac was discovered by simply including this compound in field screening tests, as in the study by Willemse *et al.* Sticky traps baited with 100 µg of Z3-14:Ac, set out in woodlands near Seewiesen (southwest of Munich, southern Germany) during June/July, caught large numbers of *C. electella*, *M. tenebrella* and another Anomologinae species, *Argolamprotes micella* (Den. & Schiff.). The three

species feed on *Picea*, *Rumex* and *Rubus*, respectively, and their relative trap captures greatly varied between test sites, depending on host abundance; which may explain why one species (*A. micella*) was missing from the test by Willemse *et al.*

The outstanding effectiveness of the Z3-14:Ac for males of these gelechiid species was supported by electroantennogram measurements. These were made from males newly taken in Z3-14:Ac baited traps (with antennae not yet glued to the adhesive), using technical procedures as in other Microlepidoptera [3, 4]. In the series of (*Z*)- and (*E*)-alkenyl acetates, varied for chain length and double bond position, the Z3-14:Ac, at the test amount of 1 µg, elicited the greatest EAG response. This was followed by the geometric isomer (*E*3-14:Ac), the corresponding alcohol analogue (Z3-14:OH) and some positional isomers and shorter-chain homologues (Z2-14:Ac, Z4-14:Ac, E4-14:Ac, Z5-14:Ac, Z3-13:Ac, Z3-12:Ac). Evaluation of equipotent stimulus amounts [3, 4] showed that these compounds were between 10- to 300-fold less stimulating than the Z3-14:Ac, in all three species. EAG effectiveness decreased by > 1,000-fold when the double bond was beyond position 5 or chain length was altered by more than 2 carbons. These data supported the view that Z3-14:Ac may be the primary female pheromone in all three species.

Further field tests were designed to assess modifying effects of second compounds. Thus, 100 µg of Z3-14:Ac and a varying amount of another compound [5] were applied to the same rubber cap dispenser. Trap types and operating procedures were as in field studies on other Microlepidoptera [4, 6]. For the three gelechiids, no attraction synergism over Z3-14:Ac alone was found but some compounds strongly reduced the response. Results of a pilot study using 100 µg additions of other compounds are presented in Table I. They show strong inhibitory effects for E3-14:Ac and Z3-14:OH in all three species and for E3-12:Ac in *C. electella*, whereas the five other additives (Z3-10:Ac, Z3-12:Ac, Z5-14:Ac, E5-14:Ac, Z5-16:Ac) did not markedly alter capture rates.

The inhibitory effects of the E3-14:Ac and Z3-14:OH were further studied in subsequent tests. Results obtained for *C. electella* and *A. micella* in two test series using 3, 10, 30 and 100 µg additions of these compounds are presented (Table II). The data show that a 30% addition of E3-14:Ac and a 3% one

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen
0341-0382/87/1100-1352 \$ 01.30/0



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

Table I. Captures of male *Chionodes electella*, *Monochroa tenebrella* and *Argolamprotes micella* in tetratrap baited with 100 µg Z3-14:Ac alone or in combination with eight different additives. Seewiesen/Starnberg, June 19 to July 27, 1982; three replicates.

Additive [100 µg]	\bar{X} males/trap of		
	<i>C. electella</i>	<i>M. tenebrella</i>	<i>A. micella</i>
none	71.0	17.7	26.7
Z3-10:Ac	61.3	22.7	22.0
Z3-12:Ac	80.3	14.0	34.0
E3-12:Ac	0	9.3	23.3
E3-14:Ac	0	0	0
Z5-14:Ac	49.7	23.0	22.0
E5-14:Ac	53.3	11.7	19.7
Z5-16:Ac	45.7	17.0	16.7
Z3-14:OH	0	0	0

Table II. Effect of varying amounts of E3-14:Ac or Z3-14:OH on captures of male *Chionodes electella* and *Argolamprotes micella* with 100 µg Z3-14:Ac. Seewiesen/Starnberg, July 3 to 29, 1983 (series A) and June 25 to July 24, 1984 (series B); tetratrap, four replicates per series.

Additive, µg	\bar{X} males/trap of			
	<i>C. electella</i>		<i>A. micella</i>	
	A	B	A	B
none	92.75	55.0	17.5	28.5
E3-14:Ac, 3	53.0	18.0	9.25	9.0
	10	18.5	9.25	2.0
	30	0.25	0	0
	100	0	0.25	—
Z3-14:OH, 3	0	0	0	1.75
	10	0.5	0	0
	30	0	0	—
	100	0	0	—

of Z3-14:OH were sufficient to suppress captures, in both species. In these tests the main flight of *M. tenebrella* had already passed when traps were set. A separate test conducted on this species (Table III) revealed inhibitory effects of the two compounds comparable to the two other species.

The abolition of captures by adding to the Z3-14:Ac source only 3% of the alcohol analogue Z3-14:OH was an unexpected discovery, as the Z3-14:Ac samples used contained almost 1% of this parent alcohol [7]. This suggests that Z3-14:Ac sources free of alcohol might be even more attractive for these gelechiids. Related alcohols, such as Z3-12:OH or E3-14:OH, showed only weak inhibitory effects.

Table III. Effect of varying amounts of E3-14:Ac or Z3-14:OH on captures of male *Monochroa tenebrella* with 100 µg Z3-14:Ac. Seewiesen/Starnberg, June 10 to July 12, 1985; tetratrap, three replicates.

Additive, µg		\bar{X} males/trap
none		37.7
E3-14:Ac, 3	10	10.7
	30	3.3
	100	0.3
	100	0
Z3-14:OH, 3	10, 30, 100	2.0
		0

On extending these tests further to various other structural analogues, E3-13:Ac was established as another strong inhibitor for *C. electella* (but not the two other species). Both E3-12:Ac and E3-13:Ac are identified sex pheromone or attractant components for some other Gelechiinae and Anomologinae [8–11]. No further strong inhibitor was found for the three test species. Particularly noteworthy is the failure of various chain length homologues (from Z3-10:Ac to Z3-18:Ac) to modify captures by Z3-14:Ac. Two of these homologues, Z3-13:Ac and Z3-12:Ac, even caught small numbers of *C. electella* and *M. tenebrella* (but not *A. micella*) when offered as single chemicals. A similar pattern of attraction specificity as for *C. electella* was obtained for two other *Chionodes* spp. responsive to Z3-14:Ac, *C. luctuella* (Hb.) and *C. nebulosella* (Hein.), at other central European test sites.

When testing binary combinations between Z3-14:Ac and another (Z)-3-alkenyl acetate, numerous male *Aproaerema anthyllidella* (Hb.) (Gelechiinae) were caught by the 100/100 mixture of Z3-14:Ac / Z3-12:Ac. The species was absent from neighbouring traps containing the single chemicals alone, indicating a synergistic attraction response. In subsequent tests of different mixtures of the two compounds, maximum attraction occurred with Z3-14:Ac / Z3-12:Ac ratios ranging from 100/10 to 100/100; the 10/100 mixture produced only poor captures and the single chemicals alone again were non-attractive (Table IV). Male *A. anthyllidella* were also caught by 100/100 combinations of Z3-14:Ac with either Z3-13:Ac or E3-12:Ac (data not shown), though at far lower numbers than with the 100/100 Z3-14:Ac / Z3-12:Ac combination. Third components were tested on this species as 100 µg amounts

Table IV. Captures of male *Aproaerema anthyllidella* in tetratraps baited with varying amounts of Z3-14:Ac and Z3-12:Ac. Seewiesen/Starnberg, July 15 to August 21, 1985; three replicates.

Amount [$\mu\text{g}/\text{trap}$] of Z3-14:Ac Z3-12:Ac		\bar{X} males/trap
100	0	0
100	3	6.7
100	10	25.7
100	30	18.3
100	100	23.7
30	100	9.0
10	100	1.3
3	100	0
0	100	0

added to this binary mixture. Only the alcohol Z3-14:OH reduced but did not fully suppress captures; various other third components tested (including Z3-10:Ac, E3-10:Ac, E3-12:Ac, Z5-12:Ac, Z3-13:Ac, E3-13:Ac, E3-14:Ac, Z5-14:Ac, Z3-16:Ac, Z5-16:Ac and Z3-12:OH) did not markedly modify capture rates. In this species, using a synergistic attractant system, attraction-inhibition thus appears to be less pronounced, compared to the other gelechiids treated above.

Potent inhibitors discovered in lepidopterous sex-attractant systems have often turned out to be essential attractant components in other, taxonomically related species. The E3-13:Ac and E3-12:Ac, highly inhibitory to male *C. electella*, are indeed the re-

ported pheromones or attractants of certain other gelechiids ([8–11]; see also above). To further test potential attractive or synergistic properties for these two compounds, and also for E3-14:Ac and Z3-14:OH, traps containing 100 μg amounts of these chemicals either alone or in different combinations with 100 μg Z3-14:Ac were set out at various central European test sites for some years. A number of further Gelechiinae and Anomologinae spp. (representing the genera *Apodia*, *Athrips*, *Bryotropha*, *Caryocolum*, *Chionodes*, *Metzneria*, *Monochroa*, *Recurvaria*, *Scrobipalpa* and *Scrobipalpopsis*), all unresponsive to pure Z3-14:Ac, were effectively captured by some of these baits. For example, male *Chionodes holosericeella* (H. Sch.) responded specifically to E3-12:Ac and male *C. perpetuella* (H. Sch.) to E3-14:Ac, with Z3-14:Ac (the *C. electella* attractant) acting inhibitorily on these species. Such data indicated a role for the various inhibitors in reproductive species isolation. Results obtained on species from the two gelechiid subfamilies responsive to $\Delta 3$ -alkenyl attractants other than Z3-14:Ac will be presented elsewhere.

Acknowledgements

I thank S. Voerman (Wageningen) and D. Reed (Saskatoon) for test chemicals; G. Dobler (Innsbruck) for conducting field tests in western Austria; and K. Burmann (Innsbruck) for help with identification of captured specimens.

- [1] H. Arn, M. Tóth, and E. Priesner, List of Sex Pheromones of Lepidoptera and Related Attractants, OILB Publ., Paris 1986.
- [2] L. P. M. Willemsse, C. J. H. Booij, and S. Voerman, J. Appl. Ent. **103**, 508 (1987).
- [3] E. Priesner, Z. Naturforsch. **38c**, 874 (1983).
- [4] E. Priesner, Z. Naturforsch. **39c**, 849 (1984).
- [5] Most chemicals were purchased from the Institute for Pesticide Research, Wageningen, The Netherlands. Additional samples were generously supplied by the Plant Biotechnology Institute, Saskatoon, Canada. All were $\geq 98\%$ pure with respect to geometric or positional isomers.
- [6] E. Priesner, H. Bogenschütz, and H. Arn, Z. Naturforsch. **35c**, 390 (1980).
- [7] S. Voerman, personal communication.
- [8] M. Renou, C. Descoins, J. Y. Lallemand, E. Priesner, M. Lettere, and M. Gallois, Z. ang. Ent. **90**, 275 (1980).
- [9] R. Baker, R. H. Herbert, N. N. Neequaye, and K. N. Rao, J. Chem. Ecol. **11**, 989 (1985).
- [10] B. F. Nesbitt, P. S. Beevor, A. Cork, D. R. Hall, R. M. Murillo, and H. R. Leal, Ent. exp. appl. **38**, 81 (1985).
- [11] D. W. Reed and M. D. Chisholm, J. Chem. Ecol. **11**, 1645 (1985).