

Pentacyclic Triterpenoid Alkanoates in Propolis

Alberto dos Santos Pereira^a, Joaquim Fernando Mendes da Silva^b, Renata Kiltzke^b,
Jari Nobrega Cardoso^a and Francisco Radler de Aquino Neto^a

^a LADETEC, Instituto de Química, Universidade Federal do Rio de Janeiro,
Ilha do Fundão, Cidade Universitária, Centro de Tecnologia, Bloco A, Lab. 607,
Rio de Janeiro, RJ – Brazil 21949–900. E-mail: ladetec@iq.ufrj.br

^b Faculdade de Farmacia, Universidade Federal do Rio de Janeiro, Ilha do Fundão, CCS,
Bloco K, 2º Andar, Rio de Janeiro, RJ – Brazil 21941–590

* Author for correspondence and reprint requests

Z. Naturforsch. **54c**, 1115–1118 (1999); received June 1/July 23, 1999

Propolis, High Temperature Gas Chromatography/MS, Triterpenyl Alkanoates,
High Molecular Weight Esters, Bee Wax

A series of high molecular mass pentacyclic terpenoid wax esters (TTFAE) were characterized for the first time in hexane extracts of Brazilian propolis (Carangola, Minas Gerais State). Identification was by HT-HRGC (High Temperature High Resolution Gas Chromatography) and HT-HRGC coupled to mass spectrometry (HT-HRGC-MS).