

Metabolites from Endophytes of the Medicinal Plant *Erythrina crista-galli*

Daniela Weber^a, Susana Gorzalczany^b, Virginia Martino^c, Cristina Acevedo^b,
Olov Sterner^{d,*}, and Timm Anke^{a,*}

^a Institut für Biotechnologie und Wirkstoff-Forschung IBWF e.V., Erwin-Schrödinger-Strasse 56, D-67663 Kaiserslautern, Germany. Fax: +496313167215.

E-mail: timm.anke@ibwf.uni-kl.de

^b Cátedra de Farmacología, Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Junín 956, 1113 Buenos Aires, Argentina

^c Cátedra de Farmacognosia, Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Junín 956, 1113 Buenos Aires, Argentina

^d Department of Organic Chemistry, University of Lund, P.O. Box 124, S-22100 Lund, Sweden. Fax: +46462228209. E-mail: Olov.Sterner@bioorganic.lth.se

* Authors for correspondence and reprint requests

Z. Naturforsch. **60c**, 467–477 (2005); received February 21/March 11, 2005

Erythrina crista-galli (Fabaceae) is used in Argentinean ethnopharmacology as anti-inflammatory medication, narcotic, disinfectant, and for the treatment of wounds. The common name of the tree is “ceibo” or coral tree. The dominating endophytes in *E. crista-galli* all belong to the genus *Phomopsis* as identified by microscopic features and the analysis of their ITS sequences. To investigate a possible contribution of *Phomopsis* spp. to the metabolites found in the plant, twelve different isolates were cultivated in different media. Besides several new metabolites a number of known compounds were detected: mellein, nectriapyrone, 4-hydroxymellein, scytalone, tyrosol, clavatol, mevinic acid, and mevalonolactone.

Key words: *Erythrina crista-galli*, Endophytic *Phomopsis*, Metabolites