Atrovirisidone B, a New Prenylated Depsidone with Cytotoxic Property from the Roots of *Garcinia atroviridis*

Dharma Permana^{a,b}, Faridah Abas^b, Maulidiani^b, Khozirah Shaari^b, Johnson Stanslas^c, Abdul Manaf Ali^d, and Nordin Hj. Lajis^{b,*}

- ^a Faculty of Medicine, Yarsi University, Jakarta 10510, Indonesia
- ^b Laboratory of Natural Products, Institute of Bioscience, University of Putra Malaysia, 43400 Serdang, Selangor, Malaysia. Fax: +60 3 89468080. E-mail: nhlajis@ibs.upm.edu.my
- ^c Faculty of Medicine and Health Sciences, University of Putra Malaysia, 43400 Serdang, Selangor, Malaysia
- ^d Laboratory of Molecular and Cells Biology, Institute of Bioscience, University of Putra Malaysia, 43400 Serdang, Selangor, Malaysia
- * Author for correspondence and reprint requests
- Z. Naturforsch. 60 c, 523–526 (2005); received February 25, 2005

A new prenylated depsidone, atrovirisidone B (2), together with naringenin (3) and 3,8"binaringenin (4) were isolated from the roots of *Garcinia atroviridis*. Their structures were determined on the basis of spectral data interpretation. Compound 2 showed cytotoxic activity against human breast (MCF-7), human prostate (DU-145) and human lung (H-460) cancer cells.

Key words: Garcinia atroviridis, Cytotoxic Activity, Atrovirisidone B