

Cytotoxic Properties of Oligostilbenoids from the Tree Barks of *Hopea dryobalanoides*

Sahidin^a, Euis H. Hakim^a, Lia D. Juliawaty^a, Yana M. Syah^a, Laily bin Din^b, Emilio L. Ghisalberti^c, Jalifah Latip^b, Ikram M. Said^b, and Sjamsul A. Achmad^{a,*}

^a Department of Chemistry, Institut Teknologi Bandung, Jalan Ganesha 10, Bandung 40132, Indonesia. E-mail: sjamsu@indo.net.id

^b School of Chemistry and Food Technology, Faculty of Science and Technology, National University of Malaysia, 43600 Bangi, Selangor, Malaysia

^c Chemistry, School of Biomedical and Chemical Sciences, The University of Western Australia, Crawley, WA 6909, Australia

* Author for correspondence and reprint requests

Z. Naturforsch. **60c**, 723–727 (2005); received April 26, 2005

A new modified stilbene dimer, diptoindonesin D (**1**), was isolated from the acetone extract of the tree bark of *Hopea dryobalanoides*, together with seven known compounds, parviflorol (**2**), (–)-balanocarpol (**3**), heimiol A (**4**), hopeafuran (**5**), (+)- α -viniferin (**6**), vaticanol B (**7**) and (–)-hopeaphenol (**8**). Cytotoxic properties of compounds **1–8** were evaluated against murine leukemia P-388 cells. Compound **8** was found to be the most active with IC₅₀ of 5.7 μ M.

Key words: Diptoindonesin D, *Hopea dryobalanoides*, Murine Leukemia P-388 Cells