

Indole Alkaloids from *Aspidosperma rigidum* and *A. schultesii* and their Antiparasitic Effects

Matías Reina^{a,*}, Wilfredo Ruiz-Mesia^a, Lastenia Ruiz-Mesia^b,
Rafael Martínez-Díaz^c, and Azucena González-Coloma^d

^a Instituto de Productos Naturales y Agrobiología (IPNA), CSIC, Avda. Astrofísico F. Sánchez 3, P. O. Box 195, 38206 – La Laguna, Tenerife, Canary Islands, Spain.
Fax: 34-922 260135. E-mail: mreina@ipna.csic.es

^b Laboratorio de Productos Naturales Antiparasitarios de la Amazonía Peruana (LIPNAA), Universidad Nacional de la Amazonía Peruana (UNAP), Iquitos, Perú

^c Departamento de Medicina Preventiva, Salud Pública y Microbiología,
Facultad de Medicina, Universidad Autónoma de Madrid (UAM), Madrid, Spain

^d Instituto de Ciencias Agrarias (ICA), CSIC, Serrano 115-dpdo, 28006 Madrid, Spain

* Author for correspondence and reprint requests

Z. Naturforsch. **66c**, 225–234 (2011); received October 7/December 21, 2010

Five oxindole alkaloids, three plumerane-type alkaloids, subtype haplophitine, and one aspidospermatane-type alkaloid, subtype tubotaiwine, were isolated from the medicinal plants *Aspidosperma rigidum* and *A. schultesii*. One compound was identified as the transoid conformer of 18-oxo-*O*-methylaspidoalbione which was not previously described. The antiparasitic activity of all compounds against *Trypanosoma cruzi* and *Leishmania infantum* and their non-specific cytotoxicity against mammalian cells were also determined.

Key words: *Aspidosperma rigidum*, *Aspidosperma schultesii*, Indole Alkaloids