

# Flavonoid Glycosides and their *p*-Coumaroyl Esters from *Campylospermum calanthum* Leaves

Savio S. Elo Manga<sup>a</sup>, Anastasie E. Tih<sup>a</sup>, Bintou Abderamane<sup>b</sup>,  
Raphael T. Ghogomu<sup>a,\*</sup>, Alain Blond<sup>c</sup>, and Bernard Bodo<sup>c</sup>

<sup>a</sup> Faculty of Science, Department of Organic Chemistry, University of Yaounde 1,  
P. O. Box 812, Yaounde, Cameroon. E-mail: ghogomut@uy1.uninet.cm

<sup>b</sup> Faculté des Sciences, Université de Ndjamen, P. O. Box 127, Ndjamen, Tchad

<sup>c</sup> Laboratoire de Chimie des Substances Naturelles, Muséum National d'Histoire  
Naturelle, 63 Rue Buffon, 75005 Paris, France

\* Author for correspondence and reprint requests

Z. Naturforsch. **67c**, 233–243 (2012); received August 19, 2011/January 20, 2012

Six new compounds, comprising three flavonoid glycosides and their respective coumaroyl esters, have been isolated and characterized from the methanol extract of the leaves of *Campylospermum calanthum*, along with three known flavonoid aglycones, 7-*O*-methyl apigenin (**1**), 7-*O*-methyl luteolin (**2**), and 7-*O*-methyl quercetin (**3**). Their structures were elucidated based on chemical evidence as well as spectroscopic analysis including 1D and 2D NMR (<sup>1</sup>H-<sup>1</sup>H COSY, HSQC, HMBC, and NOESY) spectroscopy and by comparing their spectral data with those reported for related compounds.

*Key words:* *Campylospermum calanthum*, Ochnaceae, Flavonoid Glycosides,  
Coumaroyl Esters