

# Norditerpenoid Alkaloids from *Delphinium flexuosum* Bieb.

Sevda Pırıldar<sup>a</sup>, Çağlayan Ünsal-Gürer<sup>a</sup>, Mine Koçyiğit<sup>b</sup>, Josef Zapp<sup>c</sup>,  
Alexandra K. Kiemer<sup>c</sup>, and Ali H. Meriçli<sup>a,\*</sup>

<sup>a</sup> Istanbul University, Faculty of Pharmacy, Department of Pharmacognosy,  
34116 Beyazıt, Istanbul, Turkey. Fax: +902124400252. E-mail: alimer@istanbul.edu.tr

<sup>b</sup> Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Botany,  
34116 Beyazıt, Istanbul, Turkey

<sup>c</sup> Saarland University, Institute of Pharmaceutical Biology, P. O. Box 151150,  
D-66041 Saarbrücken, Germany

\* Author for correspondence and reprint requests

Z. Naturforsch. **67c**, 541–544 (2012); received November 30, 2011/March 29, 2012

Delphiniflexine, a new norditerpenoid alkaloid, together with the three known norditerpenoid alkaloids methyllycaconitine, ajadine, and acoseptrigenine were isolated from the aerial parts of *Delphinium flexuosum*. The structure of delphiniflexine was established on the basis of <sup>1</sup>H, <sup>13</sup>C, DEPT, homonuclear <sup>1</sup>H COSY, NOESY, HSQC, and HMBC NMR studies.

*Key words:* Norditerpenoid Alkaloids, *Delphinium flexuosum*, Delphiniflexine