

Novel Betaines of the Hexaalkylguanidinio-carboxylate Type

Matthias Walter and Gerhard Maas

Institute for Organic Chemistry I, University of Ulm, Albert-Einstein-Allee 11, 89081 Ulm, Germany

Reprint requests to Prof. Dr. Gerhard Maas. Fax: 49 (0)731-5022803.

E-mail: gerhard.maas@uni-ulm.de

Z. Naturforsch. **2009**, *64b*, 1617–1624; received September 29, 2009

Dedicated to Professor Hubert Schmidbaur on the occasion of his 75th birthday

Betaines **7a**, **b** and **8a**, **b** have been prepared from 3- and 4-piperidinecarboxylic acid and *N,N,N',N'*-tetraalkyl-chloroformamidinium chlorides *via* the corresponding methyl esters. These betaines are highly hygroscopic, thermally very stable, and, with the exception of **7b**, have rather low melting points. They undergo a surprisingly facile alkaline cleavage of the hexaalkylguanidinium moiety. They react with dichloromethane by a twofold nucleophilic substitution to form methylene dicarboxylates such as **11**. The NMR (¹H, ¹³C) data of betaines **7** and **8** are discussed.

Key words: Betaines, Guanidinium, Nipecotic Acid, Isonipecotic Acid, Piperidinecarboxylic Acids

Introduction

Betaines are zwitterionic compounds containing a positively charged onium group, which does not bear a hydrogen atom, and a negatively charged functional group not adjacent to the cationic function. They have been named after (trimethylammonio)acetate (“betaine”, nowadays called glycine betaine) (Fig. 1, **A**), a common naturally occurring compound which is found in significant amounts in, *e. g.*, sugar beet, sugar beet molasses, and in shellfish [1]. Glycine betaine and related betaines have various functions in biological systems; for example, they are organic osmolytes, which protect cells against osmotic stress and dehydration [2], and act as methyl donors. Glycine betaine and other *N*-peralkylated glycinates are among the so-called compatible solutes, which can enhance the efficiency of the polymerase chain reaction by destabilization of the DNA duplex and facilitation of the initial strand dissociation process [2, 3].

Due to their ampholytic character and the resulting physico-chemical properties, the low toxicity, mild-on-skin properties and biodegradability, natural betaine **A** and synthetic betaines such as cocamidopropyl betaine (**B**) find wide application in shampoos, hand soaps, hair conditioners, and cosmetic formulations [4, 5]. Synthetic betaines, such as **B** and analogous sulfobetaines, are amphotenside ingredients in washing powders [6].

Given our interest in the properties of hexaalkylguanidinium salts, in particular those which are ionic liquids [7], we became attracted to the so far unknown betaines of the hexaalkylguanidinio-carboxylate type. When designed appropriately, such compounds could have properties similar to those of hexaalkylguanidinium-based ionic liquids, such as low melting point, high thermal stability and low vapor pressure. In addition, the zwitterionic character of the molecule and the presence of the carboxylate function in particular could endow these compounds with a reactivity which is reminiscent of task-specific ionic liquids. A number of betaines, mainly with imidazolium, tetraalkylammonium and pyridinium as the cationic group and sulfonate or carboxylate as the anionic site, have recently been studied as a new class of ion-conductive matrices for electrochemical devices [8, 9]. As target molecules, we chose the hexaalkylguanidinio-carboxylates **C** and **D**, which represent guanylated derivatives of the pharmaceutically relevant amino acids 3- and 4-piperidinecarboxylic acid (nipecotic and isonipecotic acid), respectively. Sha and Liebscher [10] have recently prepared a range of hexaalkylguanidinium salts, which have the nitrogen atom of α -amino acid esters incorporated into the guanidinium group; however, they did not proceed further to generate betaines by converting the carboxylic ester into a carboxylate function.





Fig. 1. Glycine betaine (**A**), cocamidopropyl betaine (**B**, $n = 11$), and novel betaines **C** and **D**.

Results and Discussion

Synthesis of guanidinio-carboxylates **7** and **8**


The synthesis of betaines **7** and **8** (Scheme 1) employs an established method for the preparation of hexasubstituted guanidinium salts, namely the reaction of a *sec*-amine with an N,N,N',N' -tetrasubstituted chloroamidinium salt, readily prepared from the corresponding urea and phosgene or oxalyl chloride and used without purification [11, 12]. Commercially available (\pm)-nipecotic acid (**1a**) and isonipecotic acid (**1b**) were first converted into the corresponding methyl esters **2a, b** by treatment with SOCl_2 /methanol, as described in the literature [13, 14]. These piperidinecarboxylic esters were then guanylated with chloroformamidinium salts **3** and **4** in the presence of triethylamine to give the guanidinium salts **5a, b** and **6a, b**, respectively. The guanidinium salts could not be isolated in pure form, because the separation from also formed triethylammonium chloride was incomplete. While this separation in other cases succeeds by neutralization of the ammonium salt with aqueous NaOH, we found that guanidinium salts **5** and **6** suffer rapid ester cleavage when exposed to 2 M aqueous NaOH to provide the desired betaines **7a, b** and **8a, b**, respectively.

Hexaalkylguanidinium salts are normally rather stable to hydroxide ions in dilute solution at r. t. [15, 16]. If at all, alkaline hydrolysis occurs at a significant rate only at elevated temperature: while the kinetics were studied for hexamethylguanidinium perchlorate at 60 °C [17], hexaalkylguanidinium salts bearing longer alkyl chains were reported to be stable to equimolar KOH at 60–80 °C [16]. Therefore, we were surprised to observe that betaines **7** and **8** underwent slow cleavage of the guanidinium function when exposed to 2 M aqueous NaOH at r. t. The scenario is shown exemplarily for betaine **7b** in Scheme 2. In agreement with the unsymmetrical constitution of the guanidinium group, hydrolytic cleavage proceeded in two



Scheme 1. Synthesis of betaines **7** and **8**.

directions, one leading to *N*-dimethylaminocarbonyl-4-piperidinecarboxylic acid (**9**) and dimethylamine, the other to tetramethylurea and aminoacid **1b**. The two urea derivatives were formed in a 77:23 ratio according to ¹H NMR spectroscopy. Fortunately, the

Scheme 2. Alkaline cleavage of betaine **7b**.

alkaline cleavage of betaines **7** and **8** is slow under the conditions for the generation of the betaines from esters **5** and **6**, so that the betaines can be obtained selectively under controlled reaction conditions.

In contrast to betaines **7** and **8**, the related guanidinium chloride **10** (Fig. 2), prepared by analogy to a literature method [12b], shows the typical behavior of simple hexaalkylguanidinium salts toward bases: it appears to be stable to an excess of dilute aqueous NaOH at r. t., but is decomposed slowly in concentrated aqueous NaOH.





Fig. 2.

Notably, betaines **7** and **8** cannot be obtained by direct reaction of piperidinecarboxylic acids **1** with the chloroformamidinium salts **3** and **4**. The reaction of **1b** with **3** or **4** in the presence of two equivalents of triethylamine gave rise to the formation of the corresponding tetraalkylurea besides triethylammonium chloride. Very likely, aminoacid **1b** adds to the chloroformamidinium salts through the carboxylate function, and the resulting addition product is then cleaved to yield the tetraalkylurea and the acid chloride of **1b**, which probably forms oligomers under the reaction conditions. The ability of chloroamidinium salts to activate carboxylic acids [12, 18] and to undergo chlorination reactions [12] is known.

Properties of betaines **7** and **8**

Betaines **7** and **8** are water-soluble and water-stable solids which are extremely hygroscopic and deliquesce within hours or days if not kept in a rigorously dry environment. For example, we observed that **7a, b** picked up one equivalent of water within 3–6 h of exposure to laboratory air and 5–6 equivalents of

Scheme 3. Reaction of betaine **7b** with dichloromethane.

water during 10 d.

In contrast to the solubility in water and acetonitrile, betaines **7** and **8** are almost insoluble in dichloromethane, but surprisingly they undergo a chemical reaction with this solvent. For example, when a suspension of betaine **7b** in CH_2Cl_2 was stirred for several hours, a part of the solid had disappeared. Analysis of the dichloromethane phase indicated the formation of a new product, which was identified as the dicationic salt **11** (Scheme 3) by its NMR data and an ESI-MS ($[\text{cation}]^{2+}\text{Cl}^-$ as the base peak). This reaction appears to be an equilibrium reaction which proceeded only to a conversion of 85 % of **7b** after 72 h (60 % conversion after 3 h). Obviously, the methylene dicarboxylate **11** results from a twofold nucleophilic substitution reaction at the CH_2Cl_2 molecule. The remarkably smooth formation of **11** may be attributed to an exceptionally high nucleophilicity of the carboxylate group in betaine **7b** due to the absence of ion pairing with a cationic counterion. Kinetic investigations have shown that CH_2Cl_2 is much more reluctant to an $\text{S}_{\text{N}}2$ reaction than dibromo- and diiodomethane [19]. Furthermore, the synthesis of methylene diacetate [20] and methylene dicarboxylates [21] usually requires much harsher reactions than in our case, *e. g.* by heating of CH_2X_2 ($\text{X} = \text{Br}, \text{I}$) with an excess of KOAc in refluxing acetic acid containing acetic anhydride [19].

Thermal behavior

Because of partial hydrolysis at elevated temperatures (see Experimental Section), an exact description

Table 1. ^{13}C chemical shifts (δ values in ppm, D_2O as solvent) for guanidinium betaines **7b** and **8b**, compared with their precursors.

	1b	2b	5b	7b	6b	8b
C-2	43.47	43.02	47.35, 47.60	47.98, 48.30	48.09, 48.14, 48.20, 48.24	48.70, 48.76, 48.82, 48.85
C-3	25.64	24.37	26.49, 27.50	27.71, 28.90	26.29, 26.40, 27.25, 27.40	27.46, 27.59, 28.62, 28.72
C-4	41.42	38.05	40.02	43.42	40.10	43.62, 43.64
COO^-	182.15			183.19		183.23, 183.27
COOCH_3		176.21, 52.65	177.32, 52.43		177.18/177.21, 52.59	
guanidinium- CN_3			162.79	162.90	163.07, 163.09	163.01, 163.07
NCH_3			39.13,	39.10,		
			39.43 (2 C),	39.41 (2 C),		
			39.78	39.92		
NCH_2CH_3					CH_3 : 11.96, 12.24 (3 C);	CH_3 : 11.91, 12.21 (3 C);
					CH_2 : 43.06, 43.79 (2 C), 44.12	CH_2 : 43.06, 43.79 (2 C), 44.12
<i>N</i> -butyl					C-1: 48.66, 49.12, 49.58, 49.76;	C-1: 48.55, 49.11, 49.38, 49.72;
					C-2: 28.81, 29.04, 29.08, 29.12;	C-2: 28.80, 29.06, 29.10 (2 C);
					C-3: 19.52, 19.59;	C-3: 19.50, 19.58;
					CH_3 : 13.02, 13.05 (3 C)	CH_3 : 12.97, 13.06 (3 C)

of the thermally induced phase changes of betaines **7** and **8** by differential scanning calorimetry could not be obtained. Qualitative data could be obtained, however, by heating the betaines under vacuum (0.01 mbar). The powdery solid of **7a** becomes sticky at about 80 °C, is transformed into a yellow-orange glass at 100 °C and into a very viscous oil at about 120 °C. On cooling, a glass is formed again around 100 °C and is stable on further cooling back to 20 °C. Dissolution of the glass in acetonitrile and evaporation of the solvent reconstitutes the powdery solid of **7a**. The symmetrical betaine **7b**, on the other hand, remains unchanged when heated to 160 °C. As expected, betaines **8a**, **b**, featuring longer alkyl chains and an unsymmetrical substitution at the guanidinium moiety, have lower melting points than betaines **7a**, **b**. Betaine **8a** forms a highly viscous orange-brownish oil already at 40 °C and a glass on cooling back to 20 °C. Betaine **8b** forms a glass at 60 °C and a highly viscous yellowish oil at about 80 °C; a glass is formed again when the temperature is lowered to 60 °C. In the first two heating cycles of a DSC measurement, a glass transition temperature (T_g) was registered at 23–30 °C for **8b**. In summary, all betaines except **7b** have lower melting points than reported for a wide range of other carboxylate and sulfonate betaines [9b].

The thermal stability of the betaines was determined by thermogravimetric analyses (TGA). Decomposition temperatures (T_{dec}), defined as the temperature of maximum decomposition rate, were registered at 283 (**7a**), 331 (**7b**), 269 (**8a**), and 300 °C (**8b**). However, noticeable mass loss occurred already at temperatures lower by 60–80 °C. The TGA curves gave no hints to a con-

trolled, stepwise thermal degradation of the betaines.

NMR spectra

For steric reasons, hexa-substituted guanidinium ions exist in a propeller-like configuration both in solution [22] and in the solid state [23]. Charge delocalization gives rise to a partial double bond character at the three C–N bonds, with rotation barriers that prevent free rotation at r. t. As a consequence, the NMR spectra usually become more complex [9, 22, 23f]. This is also the case for betaines **7** and **8** and their precursors, methyl esters **5** and **6**. An illustration is given in Table 1, where the ^{13}C chemical shifts of the 4-substituted betaines **7b** and **8b** and methyl esters **5b** and **6b** are compared with those of the piperidine-4-carboxylates **1b** and **2b**. It can be seen that all four *N*-methyl groups in the guanidinium systems **5b** and **7b** are magnetically non-equivalent, and that the ring carbon atoms C-2/6 as well as C-3/5 are diastereotopic. The suggested structures of **5** and **7** contain a piperidine ring in chair conformation (which is confirmed by analysis of the H,H coupling constants in the ^1H NMR spectra, see Experimental Section) and an attached tetraalkylamidinium moiety the N–C–N plane




Fig. 3. Suggested structure of guanidinium systems **5** and **7** (Newman projection along C-2(6)–C-3(5) bonds).

of which is permanently twisted against the C–N–C plane of the piperidine ring (Fig. 3). For the unsymmetrically substituted guanidinium systems **6b** and **8b**, two orientations of the amidinium moiety $R^1_2N-C-NR^2_2$ relative to the 4-substituted piperidine ring are possible. The additional isomer gives rise to a second set of signals in the NMR spectra. While the two signal sets can be distinguished in the ^{13}C spectra (*e. g.*, C-2, C-3, and most *N*-ethyl and *N*-butyl carbon atoms give rise to four signals), the 1H NMR spectra can no longer be analyzed in detail due to extensive overlap of the multiplet signals. Analogous considerations apply for the derivatives of piperidine-3-carboxylic acids, **5a**–**8a**.

Conclusion

Hexaalkylguanidinio-carboxylates **7** and **8** represent a new type of betaines. Three of them (**7a**, **8a**, **8b**) show markedly lower melting points than many known betaines with imidazolium, ammonium or pyridinium as the cationic moiety and carboxylate or sulfonate as the anionic part. With a solid→liquid phase transition below 100 °C, betaines **8a**, **b** fulfill the definition of ionic liquids (“zwitterionic ionic liquids” [9b,d]). Two remarkable chemical properties of the new betaines have been observed. One is the unexpectedly facile alkaline cleavage of the guanidinium moiety, the other one is the high nucleophilicity of the carboxylate function which allows the formation of a methylene dicarboxylate by twofold nucleophilic substitution of dichloromethane.

Experimental Section

General information

NMR spectra were recorded using a Bruker DRX 400 spectrometer (1H : 400.13 MHz, ^{13}C : 100.61 MHz). 1H NMR spectra were referenced to the residual proton signal of the solvent [$\delta_H(D_2O) = 4.79$ ppm]; m_c = centered multiplet. Signal assignments were based on H,H (COSY, TOCSY) and C,H correlation spectra (HSQC, HMBC). Mass spectra (ESI): Micromass UK, ZMD.

Piperidinecarboxylic esters (\pm)-**2a** [13] and **2b** [14] were prepared by published procedures. Acetonitrile solutions of chloroformamidinium chlorides **3** and **4** were supplied by Prof. W. Kantlehner (Hochschule Aalen).

1-[Bis(dimethylamino)methylene]-3-(methoxycarbonyl)-piperidinium chloride (**5a**); typical procedure

Methyl piperidine-3-carboxylate hydrochloride (**2a**, 0.83 g, 4.6 mmol) and a solution of *N,N,N',N'*-tetramethyl-

chloroamidinium chloride (**3**) in acetonitrile (0.91 mol/kg, 5.83 g, 5.3 mmol of **3**) were placed in a round-bottom flask equipped with a gas inlet and flushed with argon. Dry triethylamine, dried over KOH (1.5 mL, 1.09 g, 10.8 mmol), was slowly added while cooling the flask in a water bath. The reaction mixture was allowed to stir for 72 h, and the precipitated triethylammonium chloride was filtered off under an argon atmosphere. Besides small amounts of **3** and tetramethylurea, the solution still contained some triethylammonium chloride, which could not be separated from product **5a**, and was therefore used without further purification in the next step.

Guanidinium salts **5b**, **6a**, and **6b** were prepared analogously.

NMR data of salt **5a**: 1H NMR (D_2O): $\delta = 1.61$ – 2.31 (several m, 4 H, 4- H_2 , 5- H_2), 2.67–3.69 (several m, 17 H, 2- H_2 , 3-H, 6- H_2 , $N(CH_3)_2$), 3.78 (s, 3 H, OCH_3). – ^{13}C NMR (D_2O): $\delta = 22.91$ (C-5), 25.70, 26.44 (C-4); 39.23, 39.53, 39.70, 39.89 (4 NCH_3); 41.80, 43.63 (C-3); 48.46, 48.62 (C-6); 49.43, 49.78 (C-2); 52.59 (OCH_3), 163.02 (CN_3), 175.80 ($COOCH_3$).

1-[Bis(dimethylamino)methylene]-4-(methoxycarbonyl)-piperidinium chloride (**5b**)

Prepared from methyl piperidine-4-carboxylate hydrochloride (**2b**, 0.90 g, 5.0 mmol), salt **3** dissolved in acetonitrile (0.91 mol/kg, 7.50 g, 6.8 mmol) and triethylamine (1.6 mL, 1.16 g, 11.5 mmol). – 1H NMR (D_2O): $\delta = 1.66$ (dddd, $^2J = 13$ Hz, $^3J = 11$, 10 and 4 Hz, 1 H, NCH_2CH^{ax}), 1.85 (dddd, $^2J = 13$ Hz, $^3J = 11$, 10 and 4 Hz, 1 H, NCH_2CH^{ax}), 2.00 (dddd, $^2J = 13$ Hz, $^3J = 4$, 4 and 3 Hz, 1 H, NCH_2CH^{eq}), 2.03 (dddd, $^2J = 13$ Hz, $^3J = 4$, 4 and 3 Hz, 1 H, NCH_2CH^{eq}), 2.73 (tt, $^3J = 10$ and 4 Hz, 1 H, ring-4-H); 2.88, 2.89, 2.90, 2.97 (4 s, each 3 H, NCH_3); 3.15 (ddd, $^2J = 13$ Hz, $^3J = 11$ and 3 Hz, 1 H, NCH^{ax}), 3.21 (ddd, $^2J = 13$ Hz, $^3J = 11$ and 3 Hz, 1 H, NCH^{ax}), 3.42 (ddd, $^2J = 13$ Hz, $^3J = 4$ and 4 Hz, 1 H, NCH^{eq}), 3.52 (ddd, $^2J = 13$ Hz, $^3J = 4$ and 4 Hz, 1 H, NCH^{eq}), 3.70 (s, 3 H, OCH_3). – ^{13}C NMR: Table 1.

1-[(Dibutylamino)(diethylamino)methylene]-3-(methoxycarbonyl)piperidinium chloride (**6a**)

Prepared from **2b** (0.90 g, 5.0 mmol), salt **4** dissolved in acetonitrile (2.60 mol/kg, 2.69 g, 7.0 mmol) and triethylamine (1.6 mL, 1.16 g, 11.5 mmol) in dry acetonitrile (10 mL). – 1H NMR (D_2O): $\delta = 0.91$ – 1.00 (m, 6 H, butyl- CH_3), 1.18– 1.28 (m, 6 H, NCH_2CH_3), 1.32– 1.42 (m, 4 H, butyl-3- H_2), 1.46– 2.32 (several m, 8 H, ring-4- and 5- H_2 , butyl-2- H_2), 2.68– 3.76 (several m, 13 H, ring- NCH_2CH , ring-6- H_2 , butyl-1- H_2 , NCH_2CH_3), 3.79 (s, 3 H, OCH_3).

1-[(Dibutylamino)(diethylamino)methylene]-4-(methoxycarbonyl)piperidinium chloride (**6b**)

Prepared from **2b** (0.90 g, 5.0 mmol), salt **4** dissolved in acetonitrile (2.60 mol/kg, 2.85 g, 7.4 mmol) and triethylamine (1.6 mL, 1.16 g, 11.5 mmol) in dry acetonitrile (10 mL). – ¹H NMR (D₂O): δ = 0.93–1.00 (2 t, 6 H, butyl-CH₃), 1.20–1.25 (2 q, 6 H, ethyl-CH₃), 1.28–1.42 (m, 4 H, butyl-3-H₂), 1.46–1.81 (m, 5 H, butyl-2-H₂, NCH₂CH^{ax}), 1.97 (m_c, 1 H, NCH₂CH^{ax}), 2.12 (m_c, 2 H, NCH₂CH^{eq}), 2.81 (m_c, 1 H, ring-4-H), 3.09–3.52 (m, 11 H, butyl-NCH₂, ethyl-NCH₂, 3 ring-NCH₂), 3.62 (m_c, 1 H, ring-NCH^{eq}), 3.79 (s, 3 H, OCH₃). – ¹³C NMR: Table 1.

1-[Bis(dimethylamino)methylene]piperidinium 3-carboxylate (7a); typical procedure

The crude solution of ester **5a** in acetonitrile (see above) was used. The solvent was evaporated at 0.01 mbar/35 °C, and the solid residue was dissolved in demineralized water (10 mL). Aqueous NaOH (2 M) was added drop by drop until **5a** was consumed completely (3.9 mL, reaction monitoring by ¹H NMR, disappearance of the ester signal at δ = 3.78 ppm). Water and triethylamine were evaporated at 0.01 mbar/40 °C, and the solid residue was dried at 0.01 mbar/60 °C during 5 h. Betaine **7a** was extracted with dry acetonitrile (10 mL), the solvent was evaporated at 0.01 mbar/20 °C, and the solid betaine was triturated with dry ether (10 mL) and dried (0.01 mbar /60 °C, 4 h): highly hygroscopic, beige powdery solid (0.81 g, 78 % yield based on **2a**). – ¹H NMR (D₂O): two species, **A**:**B** = 6:4, δ = 1.55–1.69 (several m, 1.0 H, ring-5-H^{ax} (**A**), ring-4-H^{ax} (**B**)), 1.76–1.95 (several m, 2.0 H, ring-5-H^{eq}, ring-4-H^{ax} (**A**), ring-5-H₂ (**B**)), 2.03–2.11 (m, 0.6 H, ring-4-H^{eq} (**A**)), 2.16–2.24 (m, 0.4 H, ring-4-H^{eq} (**B**)), 2.42 (m, 0.4 H, ring-3-H (**B**)), 2.66 (m, 0.6 H, ring-3-H (**A**)), 2.93–3.10 (several s, 12.0 H, NCH₃ (**A**, **B**)), 3.10–3.24 (several m, 0.8 H, ring-2-H^{ax} and -6-H^{ax} (**B**)), 3.24–3.40 (several m, 1.8 H, ring-2-H^{ax}, 6-H₂ (**A**)), 3.51–3.60 (several m, 1.4 H, ring-2-H^{eq} (**A**), ring-2-H^{eq}, -6-H^{eq} (**B**)). – ¹³C NMR (D₂O): two species **A** and **B**, δ = 23.45 (ring-C-5 (**B**)); 23.98 (ring-C-5 (**A**)); 27.16 (ring-C-4 (**A**)); 27.87 (ring-C-4 (**B**)); 39.12, 39.25, 39.49, 39.69, 40.00 (NCH₃ (**A**, **B**)); 43.18 (ring-C-3 (**A**)), 45.28 (ring-C-3 (**B**)), 48.64 (ring-C-6 (**B**)), 48.89 (ring-C-6 (**A**)), 51.06 (ring-C-2 (**A**)), 51.41 (ring-C-2 (**B**)), 162.94 (CN₃ (**A**, **B**)), 181.41 (COO (**A**, **B**)). – MS (ESI): *m/z* (%) = 228.08 (100) [M+H]⁺.

1-[Bis(dimethylamino)methylene]piperidinium 4-carboxylate (7b)

Prepared from the crude solution of **5b** in acetonitrile as described for **7a**. Very hygroscopic, colorless powdery solid (0.87 g, 76 % yield based on **2b**). – ¹H NMR (D₂O): δ = 1.63 (dddd, ²*J* = 13.5 Hz, ³*J* = 11, 10 and 4 Hz, 1 H, 3-H^{ax}), 1.85 (dddd, ²*J* = 13.5 Hz, ³*J* = 11, 10 and 4 Hz, 1 H, 5-H^{ax}), 1.98 (dddd, ²*J* = 13.5 Hz, ³*J* = 4, 4 and 3 Hz, 1 H, 3-H^{eq}), 2.02

(dddd, ²*J* = 13.5 Hz, ³*J* = 4, 4 and 3 Hz, 1 H, 3-H^{eq}), 2.50 (tt, ³*J* = 10 and 4 Hz, 1 H, 4-H), 2.94 (s, 3 H, NCH₃), 2.96 (s, 3 H, NCH₃), 2.97 (s, 3 H, NCH₃), 3.05 (s, 3 H, NCH₃), 3.18 (ddd, ²*J* = 13 Hz, ³*J* = 11 and 3 Hz, 1 H, NCH^{ax}), 3.25 (ddd, ²*J* = 13 Hz, ³*J* = 11 and 3 Hz, 1 H, NCH^{ax}), 3.48 (ddd, ²*J* = 13 Hz, ³*J* = 4 and 4 Hz, 1 H, NCH^{eq}), 3.58 (ddd, ²*J* = 13 Hz, ³*J* = 4 and 4 Hz, 1 H, NCH^{eq}). – ¹³C NMR: Table 1.

1-[(Dibutylamino)(diethylamino)methylene]piperidinium 3-carboxylate (8a)

Prepared from the crude solution of **6a** in acetonitrile as described for **7a**. Very hygroscopic, beige powdery solid (1.35 g, 80 % yield based on **2a**). – ¹H NMR (D₂O): two species, **A**:**B** = 6:4, δ = 0.94–0.99 (m, 6 H, butyl-CH₃ (**A**, **B**)), 1.18–1.25 (m, 6 H, ethyl-CH₃ (**A**, **B**)), 1.26–1.42 (m, 4 H, butyl-3-H₂ (**A**, **B**)), 1.44–1.82 (several m, 5.4 H, ring-4-H^{ax} (**A**), ring-4-H^{ax}, -5-H^{ax} (**B**), butyl-2-H₂ (**A**, **B**)), 1.82–1.97 (several m, 1.6 H, ring-5-H₂ (**A**), ring-5-H^{eq} (**B**)), 2.08 (d, 0.6 H, ring-4-H^{eq} (**A**)), 2.21 (d, 0.4 H, ring-4-H^{eq} (**B**)), 2.41 (m, 0.4 H, ring-NCH₂CH (**B**)), 2.65 (m, 0.6 H, ring-NCH₂CH (**A**)), 3.08–3.64 (several m, 12 H, ring-2-H₂, 6-H₂ (**A**, **B**), butyl-NCH₂ (**A**, **B**), NCH₂CH₃ (**A**, **B**)). – ¹³C NMR (D₂O): two species **A** and **B**, δ = 11.84, 11.95, 12.08, 12.14, 12.17, 12.22 (ethyl-CH₃ (**A**, **B**)); 12.97, 13.03, 13.07 (butyl-CH₃ (**A**, **B**)); 19.48, 19.54, 19.57 (butyl-C-3 (**A**, **B**)); 23.14, 23.27 (ring-C-5 (**B**)); 23.77, 23.85 (ring-C-5 (**A**)); 27.08, 27.17 (ring-C-4 (**A**)); 27.85, 27.94 (ring-C-4 (**B**)); 28.75, 28.85, 28.95, 28.99, 29.01, 29.10, 29.18 (butyl-C-2 (**A**, **B**)); 42.76, 42.88 (ring-C-3 (**A**)); 43.04, 43.09, 43.63, 43.73, 43.83, 43.86, 43.91, 44.15 (NCH₂CH₃ (**A**, **B**)); 44.86, 44.90 (ring-C-3 (**B**)); 48.56, 48.70, 49.00, 49.15, 49.30, 49.41, 49.47, 49.58, 49.69 (ring-C-6 (**A**, **B**); butyl-NCH₂ (**A**, **B**)); 51.52, 51.72, 51.78, 51.80 (ring-C-2 (**A**, **B**)); 163.11, 163.16 (CN₃ (**A**, **B**)); 180.77, 180.87, 181.24 (COO (**A**, **B**)).

1-[(Dibutylamino)(diethylamino)methylene]piperidinium 4-carboxylate (8b)

Prepared from the crude solution of **6b** in acetonitrile as described for **7a**. Very hygroscopic, beige powdery solid (1.39 g, 82 % based on **2b**). – ¹H NMR (D₂O): δ = 0.94–0.99 (m, 6 H, butyl-CH₃), 1.18–1.25 (m, 6 H, ethyl-CH₃), 1.30–1.42 (m, 4 H, butyl-3-H₂), 1.45–1.78 (m, 5 H, butyl-2-H₂, NCH₂CH^{ax}), 1.86–1.95 (m, 1 H, NCH₂CH^{ax}), 1.98–2.09 (m, 2 H, NCH₂CH^{eq}), 2.47–2.53 (m, 1 H, ring-4-H), 3.11–3.52 (m, 11 H, butyl-NCH₂, ethyl-NCH₂, 3 ring-NCH), 3.60–3.66 (m, 1 H, ring-NCH). – ¹³C NMR: Table 1.

Alkaline hydrolysis of betaine 7b

Aqueous NaOH (2 M, 2 mL) was added to a solution of betaine **7b** (0.253 g, 1.11 mmol) in demineralized water

(5 mL). After stirring for 1 h, the solvent was evaporated at 0.01 mbar/40 °C. The solid residue was triturated with dry ether (10 mL) to remove residual tetramethylurea and dissolved in water. After neutralization with dilute aqueous HCl, the solution was evaporated to dryness (0.01 mbar/80 °C, 3 h). According to the ¹H NMR spectra, the solid contained 1-(dimethylcarbamoyl)piperidine-4-carboxylic acid (**9**) [24] and piperidine-4-carboxylic acid (**1b**) in a 77:23 ratio. – NMR data of **9**: ¹H NMR (D₂O): δ = 1.51–1.61 and 1.83–1.92 (2 m_c, each 2 H, 3,5-H₂), 2.52–2.57 (m_c, 1 H, 4-CH), 2.78 (s, 3H, NCH₃), 2.79 (s, 3 H, NCH₃), 2.89 (ddd, 2 H, NCH^{ax}), 3.68 (dt, 2 H, NCH^{eq}). – ¹³C NMR (D₂O): δ = 28.90 (C-3,5), 38.09 (N(CH₃)₂), 44.57 (C-4), 46.44 (C-2,6), 166.02 (CN₃), 184.48 (COO).

1-[Bis(dimethylamino)methylene]piperidinium chloride (**10**)

Under an argon atmosphere, dry piperidine (0.8 mL, 0.69 g, 8.1 mmol) was added to a solution of salt **3** in acetonitrile (6.6 mL, 0.91 mol/kg, 5.50 g, 5.0 mmol). After stirring for 20 h, the solvent was evaporated at 0.01 mbar/30 °C, and the residue was dissolved in demineralized water (5 mL). After neutralization of piperidinium chloride with 2 M aqueous NaOH (4 mL), water and piperidine were evaporated at 0.01 mbar/40 °C. The solid residue was triturated with 2×5 mL pentane and dried (0.01 mbar/80 °C, 5 h). Very hygroscopic, colorless powdery solid (0.79 g, 89 % yield based on piperidine), m. p. 173–175 °C. – ¹H NMR (D₂O): δ = 1.52–1.76 (several m, 6 H, 3-, 4-, 5-H₂), 2.87 (s, 6 H, N(CH₃)₂), 2.91 (s, 6 H, N(CH₃)₂), 3.14–3.32 (m_c, 2-, 6-H₂). – ¹³C NMR (D₂O): δ = 23.19 (C-4), 24.83 (C-3,5); 39.24, 39.38 (N(CH₃)₂); 49.28 (C-2,6), 162.80 (CN₃).

4,4'-[Methylenebis(oxycarbonyl)]-bis{1-[bis(dimethylamino)methylene]piperidinium} dichloride (**11**)

A suspension of betaine **7b** (0.455 g, 2.0 mmol) in dry dichloromethane (20 mL) was stirred for 72 h. At this point, most of the betaine was consumed, and further reaction appeared not to take place. The residual betaine was removed by filtration under an argon atmosphere, and the solution was

evaporated to dryness at 0.01 mbar/20 °C to leave a very hygroscopic, colorless powdery solid (0.398 g), which consisted of dicationic salt **11** contaminated with about 8–10 % of betaine **7b**. Due to similar solubilities, complete separation of the two components was not possible, but a small sample enriched in **11** could be obtained by repeated trituration with CH₂Cl₂. Spectroscopic data of **11**: ¹H NMR (D₂O): δ = 1.74 (m_c, 2 H, NCH₂CH^{ax}), 1.93 (m_c, 2 H, NCH₂CH^{ax}), 2.12 (m_c, 4 H, NCH₂CH^{eq}), 2.87 (m_c, 2 H, ring-4-H); 2.96, 2.97, 2.98 and 3.04 (4 s, each 6 H, NCH₃); 3.27 (m_c, 4 H, NCH^{ax}), 3.50 (m_c, 2 H, NCH^{eq}), 3.60 (m_c, 2 H, NCH^{eq}), 5.90 (s, 2 H, OCH₂O). – ¹³C NMR (D₂O): δ = 26.36, 27.33 (C-3,5); 39.18, 39.51 (N(CH₃)₂); 39.85 (C-4); 47.33, 47.52 (C-2,6); 80.42 (OCH₂O), 162.92 (CN₃), 175.05 (COO). – MS (ESI): *m/z* (%) = 505.36 (33) [cation]²⁺ + ³⁷Cl⁻, 503.33 (100) [cation]²⁺ + ³⁵Cl⁻, 453.36 (13), 423.36 (5).

Thermal analysis

Efforts to characterize the thermal behavior of betaines **7** and **8** by differential scanning calorimetry (DSC) were hampered by a partial hydrolysis at elevated temperatures. For example, betaine **7b** was submitted to three heating cycles up to 250 °C with a heating rate of 10 °C min⁻¹. A ¹H NMR spectrum of the sample then indicated the presence of betaine **7b** (75 %), acid **9** (22 %) and tetramethylurea (3 %). Therefore, qualitative observations were made on betaine samples which were kept in a Schlenk flask under vacuum (0.01 mbar) and heated with an external oil bath. The thermal stability of the salts was determined by thermogravimetric analysis using a Mettler-Toledo TGA/SDTA 851 instrument. The temperature was increased linearly at 10 °C min⁻¹ under nitrogen from 25 to 800 °C.

Acknowledgements

This work was supported by a grant from the Bundesministerium für Bildung und Forschung (BMBF). We thank Prof. Willi Kantlehner (Hochschule Aalen) for samples of chloroformamidinium salts.

-
- [1] F.J. de Zwart, S. Slow, R.J. Payne, M. Lever, P.M. George, J.A. Gerrard, S.T. Chambers, *Food Chem.* **2003**, *83*, 197–204.
- [2] a) L. N. Csonka, *Microbiol. Rev.* **1989**, *53*, 121–147; b) M. Ashraf, M. R. Foolad, *Environ. Exp. Bot.* **2006**, *59*, 206–216; c) T. H. H. Chen, N. Murata, *Trends in Plant Science* **2008**, *13*, 499–505; d) J. M. Wood, *Microbiol. Mol. Biol. Rev.* **1999**, *63*, 230–262.
- [3] a) W. Henke, K. Herdel, K. Jung, D. Schnorr, S. A. Loening, *Nucleic Acids Res.* **1997**, *25*, 3957–3958; b) M. Ralser, R. Querfurth, H.-J. Warnatz, H. Lehrach, M.-L. Yaspo, S. Krobitsch, *Biochem. Biophys. Res. Commun.* **2006**, *347*, 747–751.
- [4] K. Koumoto, H. Ochiai, N. Sugimoto, *Tetrahedron* **2008**, *64*, 168–174.
- [5] L. Rigano, G. Dell'Acqua, R. Leporatti, *Cosmetics & Toiletries* **2000**, *115*, 47–54.
- [6] E. G. Lomax (ed.), *Amphoteric Surfactants*, 2nd ed., Surfactant Science Series 59, Marcel Dekker, New York, **1996**.

- [7] H. Kunkel, G. Maas, *Eur. J. Org. Chem.* **2007**, 3746–3757.
- [8] a) J. Cardoso, A. Huanosta, O. Manero, *Macromolecules* **1991**, *24*, 2890–2895; b) M. Galin, A. Mathis, J.-C. Galin, *Polym. Adv. Technol.* **2001**, *12*, 574–582, and refs. cited therein.
- [9] a) M. Yoshizawa, M. Hirao, K. Ito-Akita, H. Ohno, *J. Mater. Chem.* **2001**, *11*, 1057–1062; b) M. Yoshizawa, A. Narita, H. Ohno, *Aust. J. Chem.* **2004**, *57*, 139–144; c) M. Yoshizawa, H. Ohno, *Chem. Lett.* **2004**, *33*, 1594–1595; d) account: H. Ohno, *Bull. Chem. Soc. Jpn.* **2006**, *79*, 1665–1680; e) J. N. Krishnan, H. S. Kim, J. K. Lee, B.-W. Cho, E. J. Roh, S.-G. Lee, *Bull. Kor. Chem. Soc.* **2008**, *29*, 1705–1710.
- [10] J. Sha, J. Liebscher, *Synthesis* **2008**, 917–920.
- [11] a) W. Kantlehner, E. Haug, W. W. Mergen, P. Speh, T. Maier, J. J. Kapassakalidis, H.-J. Bräuner, *Synthesis* **1983**, 904–905; b) W. Kantlehner, E. Haug, W. W. Mergen, P. Speh, J. J. Kapassakalidis, H.-J. Bräuner, H. Hagen, *Liebigs Ann. Chem.* **1984**, 108–126.
- [12] T. Isobe, T. Ishikawa, *J. Org. Chem.* **1999**, *64*, 6984–6988.
- [13] N. Tamura, Y. Matsushita, T. Iwama, S. Harada, S. Kishimoto, K. Itoh, *Chem. Pharm. Bull.* **1991**, *39*, 1199–1212.
- [14] J. Alfaro-Lopez, T. Okayama, K. Hosohata, P. Davis, F. Porreca, H. Yamamura, V. J. Hruby, *J. Med. Chem.* **1999**, *42*, 5359–5368.
- [15] H. Lecher, F. Graf, *Liebigs Ann. Chem.* **1924**, 438, 154–169.
- [16] a) N. M. M. Mateus, L. C. Branco, M. M. T. Lourenço, C. A. M. Afonso, *Green Chem.* **2003**, *5*, 347–352; b) P. S. Kulkarni, L. C. Branco, J. G. Crespo, M. C. Nunes, A. Raymundo, C. A. M. Afonso, *Chem. Eur. J.* **2007**, *13*, 8478–8488.
- [17] R. B. Homer, K. W. Alwis, *J. Chem. Soc., Perkin Trans. 2* **1976**, 781–784.
- [18] H.-B. König, W. Schröck, K.-G. Metzger, H. Disselkötter, Ger. Offen. DE2104579 (A1), **1972**.
- [19] J. Hine, S. J. Ehrenson, W. H. Brader, *J. Am. Chem. Soc.* **1956**, *78*, 2282–2284.
- [20] J. G. Atkinson, D. W. Cillis, R. S. Stuart, *Can. J. Chem.* **1969**, *47*, 477–479.
- [21] K. T. Sloan, US 19630326, **1965**.
- [22] a) H. Kessler, D. Leibfritz, *Chem. Ber.* **1971**, *104*, 2158–2169; b) A. V. Santoro, G. Mickevicius, *J. Org. Chem.* **1979**, *44*, 117–120.
- [23] Hexamethylguanidinium ion: a) X. Zhang, R. Bau, J. A. Sheehy, K. O. Christe, *J. Fluorine Chem.* **1999**, *98*, 121–126; b) A. A. Kolomeitsev, G. Bissky, J. Barten, N. Kalinovich, E. Lork, G.-V. Rösenthaller, *Inorg. Chem.* **2002**, *41*, 6118–6124; c) G.-V. Rösenthaller, E. Lork, G. Bissky, A. A. Kolomeitsev, *Z. Krist. – New Cryst. Struct.* **2002**, *217*, 419–420; d) W. Frey, M. Vettel, K. Edelmann, W. Kantlehner, *Z. Krist. – New Cryst. Struct.* **1998**, *213*, 77–78. For selected other hexa-substituted guanidinium ions, see: e) W. Nakanishi, A. Ogino, M. Kawahata, K. Yamaguchi, T. Ishikawa, *Tetrahedron Lett.* **2007**, *48*, 8526–8530; f) M. Tanaka, H.-U. Siehl, T. Viefhaus, W. Frey, W. Kantlehner, *Z. Naturforsch.* **2009**, *64b*, 765–772.
- [24] A synthesis of the methyl ester of **9** from methyl piperidine-4-carboxylate and *N,N*-dimethylcarbamoyl chloride has been reported: H. L. Yale, K. Losee, J. Martins, M. Holsing, F. M. Perry, J. Bernstein, *J. Am. Chem. Soc.* **1953**, *75*, 1933–1942.