

# Synthesis and Structural Characterization of the Dinuclear Beryllium Species $[\text{Be}_2\text{Cl}_2(\mu\text{-Cl})_2(\text{PCy}_3)_2]$

Holger Braunschweig and Katrin Größ

Institut für Anorganische Chemie, Julius-Maximilians-Universität Würzburg,  
Am Hubland, 97074 Würzburg, Germany

Reprint requests to Prof. Dr. Holger Braunschweig. Fax: (+49) (0)931 / 888-4623.  
E-mail: h.braunschweig@mail.uni-wuerzburg.de

*Z. Naturforsch.* **2011**, *66b*, 55–57; received October 6, 2010

The synthesis and full characterization of  $[\text{Be}_2\text{Cl}_2(\mu\text{-Cl})_2(\text{PCy}_3)_2]$ , which results from the reaction of  $[\text{Pd}(\text{PCy}_3)_2]$  and  $\text{BeCl}_2$  with concomitant precipitation of elemental palladium, is reported.

**Key words:** Beryllium, Dinuclear Compound, Lewis Base, Palladium, X-Ray Diffraction

## Introduction


Due to the toxicity of beryllium compounds, the chemistry of beryllium is far less developed than that of its neighboring elements [1, 2]. As beryllium-containing materials feature unique properties, most of the corresponding research is done in material sciences [3]. With regard to its toxicity, additional work is focused on the coordination chemistry of Be(II) in aqueous solutions [4, 5]. Thus, tetrahedral four-coordinate Be species, which result from the coordination of ligands containing main group substituents, are well established [6–10]. However, the chemistry of transition metal-beryllium interactions was limited to cluster compounds, particularly to examples consisting of Zr [11, 12].

Based on previous success with the facile formation of “metal-only” Lewis pairs between electron-rich  $\text{Pt}^0$  complexes and *p*-block metals, *e. g.* in the case of  $[(\text{Cy}_3\text{P})_2\text{Pt}-\text{AlCl}_3]$  [13] and  $[(\text{Cy}_3\text{P})_2\text{Pt}-\text{GaCl}_3]$  [14], we sought to extend this rather unusual bonding pattern to Lewis-base adducts between *d*- and *s*-block metals.  $\text{BeCl}_2$  as a strong Lewis acid proved to be a promising starting material, and reaction of  $[\text{Pt}(\text{PCy}_3)_2]$  with  $\text{BeCl}_2$  in benzene resulted in the platinum beryllium adduct  $[(\text{Cy}_3\text{P})_2\text{Pt}-\text{BeCl}_2]$  (**1**) comprising an unprecedented, electron precise bond between beryllium and a *d*-block metal [15]. Recent studies showed that related low-valent palladium complexes also show a propensity to act as metal bases towards metal-coordinate boryl and borylene ligands, and therefore behave similar to their platinum congeners [16–19]. In the present paper we report on the reaction of  $[\text{Pd}(\text{PCy}_3)_2]$  with

$\text{BeCl}_2$ , resulting in the formation of a dinuclear beryllium species on a different reaction pathway.

## Results and Discussion

The reaction of  $[\text{Pd}(\text{PCy}_3)_2]$  and  $\text{BeCl}_2$  was conducted under similar conditions as applied to the synthesis of **1**. Thus, a toluene solution of the palladium complex was treated with a slight excess of  $\text{BeCl}_2$  and heated to 80 °C. The reaction was monitored by  $^{31}\text{P}$  NMR spectroscopy, revealing a new signal at 33.3 ppm, which is slightly highfield shifted with regard to that of the starting material (39.2 ppm). Completion of the conversion, though, required additional  $\text{BeCl}_2$  – two equivalents in total – and extended heating to 80 °C, after which a grey solid precipitated from the yellow-green solution, which we assumed to be elemental palladium.  $^9\text{Be}$  NMR spectroscopy of the new compound gave a broad signal at 12.7 ppm, which is comparable to three-coordinate beryllium compounds such as  $\text{ArBeCl}(\text{OEt}_2)$  ( $\text{Ar} = \text{C}_6\text{H}_3\text{-2,6-Me}_2$ ) (12.8 ppm), and thus more deshielded than corresponding adducts comprising beryllium in coordination number four, *e. g.*  $\text{BeCl}_2(\text{OEt}_2)_2$  and  $\text{Be}\{[\text{N}(\text{SiMe}_3)]_2\text{CPh}\}_2$ , which exhibit resonances at 2.6 and 5.5 ppm, respectively [20]. It should be noted though, that in case of the corresponding Pt-Be adduct **1** no  $^9\text{Be}$  NMR signal could be detected due to unresolved coupling to platinum and phosphorus nuclei. The aforementioned spectroscopic data as well as the required twofold excess of  $\text{BeCl}_2$  already indicate that  $[\text{Pd}(\text{PCy}_3)_2]$  does not form a Lewis base adduct with  $\text{BeCl}_2$  in analogy to its plat-



inum congener. Indeed, after work up, 62 % of a colorless, crystalline material was isolated. X-Ray analysis revealed the formation of the dinuclear phosphine adduct  $[\text{Be}_2\text{Cl}_2(\mu\text{-Cl})_2(\text{PCy}_3)_2]$  (**2**).

In the crystal, **2** displays  $C_{2h}$  symmetry, and each beryllium center is surrounded by a phosphine group, two bridging and one terminal chloride substituent, thus exhibiting a distorted tetrahedral geometry (Fig. 1).

The Be–P bond length in **2** (1.932(2) Å) is significantly shorter than in a related bisphosphine complex of beryllium  $[\text{BeCl}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)_2]$  (2.206(3) Å), recently reported by Dehnicke *et al.* [21]. The P–Be–Cl1 plane in **2** is orientated almost perpendicular with respect to the central Be–Cl2–Be<sub>a</sub>–Cl2<sub>a</sub> plane (88.2°). As to be expected, the exocyclic Be–Cl1 distance of 1.932(2) Å is shorter than the endocyclic Be–Cl separations (Be–Cl2 2.088(2) Å). Comparison to the structurally related  $(\text{Ph}_4\text{P})_2[\text{Be}_2\text{Cl}_6]$  reveals many similarities [22]. Thus, the terminal (1.952(3) Å) and the bridging Be–Cl bonds (2.102(3) Å) in the latter species are only slightly longer than those in **2**. Likewise, the central four-membered ring in **2** displays an angle of 97.13(9)° (Cl2–Be–Cl2<sub>a</sub>), which resembles




Fig. 1. Molecular structure of  $[\text{Be}_2\text{Cl}_2(\mu\text{-Cl})_2(\text{PCy}_3)_2]$  (**2**). Displacement ellipsoids are at the 50 % probability level. Symmetry related positions ( $-x+1, -y, -z+1$ ) are marked with <sub>a</sub>. Selected bond lengths (Å) and angles (deg): Be–Cl1 1.932(2), Be–P 2.216(2), Be–Cl2 2.088(2); P–Be–Cl1 114.47(10), P–Be–Cl2 110.54(10), Cl2–Be–Cl2<sub>a</sub> 97.13(9).

that in  $[\text{Be}_2\text{Cl}_6]^{2-}$  (95.6(1)°). Somewhat surprising, a CCDC search provided no information as to other structurally characterized beryllium chloride species of the type *trans*-(L)ClBe-( $\mu\text{-Cl}$ )<sub>2</sub>-BeCl(L) (L = neutral donor), wherein one donating ligand stabilizes a tetrahedral beryllium center. However, similar structural motifs are known from *d*-block metal species such as the corresponding dinuclear mercury compound  $[\text{Hg}_2(\text{Cl})_2(\mu\text{-Cl})_2(\text{PCy}_3)_2]$  [23] and the palladium complex  $[\text{Pd}_2(\text{Cl})_2(\mu\text{-Cl})_2(\text{PCy}_3)_2]$  [24]. However, the latter compound displays the expected square-planar geometry at the palladium centers.

In conclusion, we have shown that the reaction of  $[\text{Pd}(\text{PCy}_3)_2]$  with  $\text{BeCl}_2$  takes a completely different course than that of the corresponding platinum phosphine species. In case of the former,  $\text{BeCl}_2$  abstracts the phosphine ligands with formation of a dinuclear, structurally rare Be–P adduct, without any indication for the formation of a palladium-beryllium complex. Presumably, this finding can be ascribed to a decreased Lewis basicity of Pd in comparison to Pt.

## Experimental Section

*Safety note:* in view of the toxicity of beryllium and its compounds, all necessary safety measures were undertaken. All reactions were carried out on a small scale, and for NMR spectroscopy we used exclusively J. Young NMR tubes. The glassware was cleaned separately, and all waste was collected in suitable containers.

*General considerations:* All manipulations were performed under an inert atmosphere of dry argon using either standard Schlenk-line or glovebox techniques. Toluene was distilled over sodium and stored over molecular sieves prior to use.  $\text{C}_6\text{D}_6$  was dried over molecular sieves and degassed by three freeze-pump-thaw cycles before use. Anhydrous  $\text{BeCl}_2$  was purchased from Aldrich,  $[\text{Pd}(\text{PCy}_3)_2]$  was prepared according to known methods [25]. The NMR spectra were recorded on a Bruker Avance 500 ( $^1\text{H}$ : 500.13 MHz;  $^{13}\text{C}$ : 125.76 MHz;  $^{31}\text{P}$ : 202.45 MHz;  $^9\text{Be}$ : 70.28 MHz) FT-NMR spectrometer.  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra were referenced to external TMS *via* the signal of the residual protons of the solvent ( $^1\text{H}$ ) or *via* the solvent itself ( $^{13}\text{C}$ ).  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra were referenced to 85 %  $\text{H}_3\text{PO}_4$ ,  $^9\text{Be}$  NMR spectra to an aqueous solution of  $\text{BeCl}_2$ .

### *Di- $\mu$ -chloro-trans-dichloro-bis[tricyclohexylphosphine]-diberyllium (2)*

A small excess of  $\text{BeCl}_2$  (2.8 mg, 0.035 mmol) was added to a pale-yellow solution of  $[\text{Pd}(\text{PCy}_3)_2]$  (20 mg, 0.030 mmol) in toluene (0.4 mL). The reaction was heated for 18 h at 80 °C. A second portion of  $\text{BeCl}_2$  (2.0 mg,

0.025 mmol) was added, and the reaction mixture was again heated for 18 h at 80 °C to complete the conversion. No ligand exchange was observed in solution. Palladium precipitated as a dark-grey solid from the yellow-green solution, and after filtration the latter was layered with hexane. After slow evaporation in a glovebox at r.t., **2** was obtained as colorless crystals (13 mg, 62%). The crystals were redissolved in C<sub>6</sub>D<sub>6</sub> for spectroscopic characterization. – <sup>1</sup>H NMR (500.13 MHz, C<sub>6</sub>D<sub>6</sub>): δ = 1.26 (br s, 18H, Cy), 1.75–1.61 (m, 30H, Cy), 2.11–2.05 (m, 18H, Cy). – <sup>13</sup>C NMR (125.76 MHz, C<sub>6</sub>D<sub>6</sub>): δ = 26.55 (s, C<sup>4</sup>, Cy), 27.81 (virtual triplet, *N* [26] = 11 Hz, C<sup>2</sup>, C<sup>6</sup>, Cy), 31.45 (s, C<sup>3</sup>, C<sup>5</sup>, Cy), 34.14 (virtual triplet, *N* [27] = 18 Hz, C<sup>1</sup>, Cy). – <sup>31</sup>P{<sup>1</sup>H} NMR (202.46 MHz, C<sub>6</sub>D<sub>6</sub>): δ = 33.25. – <sup>9</sup>Be NMR (70.28 MHz, C<sub>6</sub>D<sub>6</sub>): δ = 12.72 (br s). – C<sub>36</sub>H<sub>66</sub>Be<sub>2</sub>Cl<sub>4</sub>P<sub>2</sub> (720.70): calcd. C 60.00, H 9.23; found C 59.19, H 8.58.

#### X-Ray structure determination

The crystal data of **2** were collected on a Bruker APEX diffractometer with CCD area detector and graphite-

monochromatized MoK<sub>α</sub> radiation. The structure was solved using Direct Methods, expanded using Fourier techniques and refined with the SHELX software package [28]. All non-hydrogen atoms were refined anisotropically. Hydrogen atoms were assigned idealized positions and were included in structure factor calculations. Crystal data for **2**: C<sub>42</sub>H<sub>72</sub>Be<sub>2</sub>Cl<sub>4</sub>P<sub>2</sub>, *M<sub>r</sub>* = 798.76, colorless block, 0.28 × 0.1 × 0.1 mm<sup>3</sup>, monoclinic space group *P*2<sub>1</sub>/*c*, *a* = 15.583(1), *b* = 8.2539(6), *c* = 18.4827(13) Å, β = 112.999(1)°, *V* = 2188.3(3) Å<sup>3</sup>, *Z* = 2, ρ<sub>calcd</sub> = 1.21 g cm<sup>-3</sup>, μ = 0.4 mm<sup>-1</sup>, *F*(000) = 860 e, *T* = 168(2) K, *R*1 = 0.0441, *wR*2 = 0.0911 for 4281 independent reflections [2θ ≤ 52.02°] and 226 refined parameters, Δρ (max / min) = 0.495 / -0.363 e Å<sup>-3</sup>.

CCDC 795511 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

#### Acknowledgement

Financial support by the DFG is gratefully acknowledged.

- [1] O. Kumberger, H. Schmidbaur, *Chem. Unserer Zeit* **1993**, *27*, 310–316.
- [2] D. N. Skilleter, *Chem. Br.* **1990**, *26*, 26–30.
- [3] A. V. G. Chizmeshya, C. J. Ritter, T. L. Groy, J. B. Tice, J. Kouvetakis, *Chem. Mater.* **2007**, *19*, 5890–5901.
- [4] H. Schmidbaur, *Coord. Chem. Rev.* **2001**, *215*, 223–242.
- [5] C. Y. Wong, J. D. Woolins, *Coord. Chem. Rev.* **1994**, *130*, 243–273.
- [6] M. P. Dressel, S. Nogai, R. J. F. Berger, H. Schmidbaur, *Z. Naturforsch.* **2003**, *58b*, 173–182.
- [7] B. Neumüller, W. Petz, K. Dehnicke, *Z. Anorg. Allg. Chem.* **2008**, *634*, 662–668.
- [8] F. Ceconi, C. A. Ghilardi, A. Ienco, P. Mariani, C. Mealli, S. Midollini, A. Orlandini, A. Vacca, *Inorg. Chem.* **2002**, *41*, 4006–4017.
- [9] K. Dehnicke, B. Neumüller, *Z. Anorg. Allg. Chem.* **2008**, *634*, 2703–2728.
- [10] B. Neumüller, K. Dehnicke, *Z. Anorg. Allg. Chem.* **2010**, *636*, 1438–1440.
- [11] R. P. Ziebarth, J. D. Corbett, *J. Am. Chem. Soc.* **1985**, *107*, 4571–4573.
- [12] J. B. Willems, H. W. Rohm, C. Geers, M. Köckerling, *Inorg. Chem.* **2007**, *46*, 6197–6203.
- [13] H. Braunschweig, K. Gruss, K. Radacki, *Angew. Chem.* **2007**, *119*, 7929–7931; *Angew. Chem. Int. Ed.* **2007**, *46*, 7782–7784.
- [14] H. Braunschweig, K. Gruss, K. Radacki, *Inorg. Chem.* **2008**, *47*, 8595–8597.
- [15] H. Braunschweig, K. Gruss, K. Radacki, *Angew. Chem.* **2009**, *121*, 4303–4305; *Angew. Chem. Int. Ed.* **2009**, *48*, 4239–4241.
- [16] H. Braunschweig, K. Radacki, D. Rais, G. R. Whittell, *Angew. Chem.* **2005**, *117*, 1217–1219; *Angew. Chem. Int. Ed.* **2005**, *44*, 1192–1194.
- [17] H. Braunschweig, D. Rais, K. Uttinger, *Angew. Chem.* **2005**, *117*, 3829–3832; *Angew. Chem. Int. Ed.* **2005**, *44*, 3763–3766.
- [18] H. Braunschweig, K. Radacki, D. Rais, F. Seeler, *Angew. Chem.* **2006**, *118*, 1087–1090; *Angew. Chem. Int. Ed.* **2006**, *45*, 1066–1069.
- [19] H. Braunschweig, C. Burschka, M. Burzler, S. Metz, K. Radacki, *Angew. Chem.* **2006**, *118*, 4458–4461; *Angew. Chem. Int. Ed.* **2006**, *45*, 4352–4355.
- [20] M. Niemeyer, P. P. Power, *Inorg. Chem.* **1997**, *36*, 4688–4696.
- [21] G. Frenking, N. Holzmann, B. Neumüller, K. Dehnicke, *Z. Anorg. Allg. Chem.* **2010**, *636*, 1772–1775.
- [22] B. Neumüller, F. Weller, K. Dehnicke, *Z. Anorg. Allg. Chem.* **2003**, *629*, 2195–2199.
- [23] L. Liu, Q. F. Zhang, W. H. Leung, *Acta Crystallogr.* **2004**, *E60*, m394–m395.
- [24] C. Sui-Seng, F. Belanger-Gariepy, D. Zargarian, *Acta Crystallogr.* **2003**, *E59*, m620–m621.
- [25] V. V. Grushin, C. Bensimon, H. Alper, *Inorg. Chem.* **1994**, *33*, 4804–4806.
- [26]  $N = |^2J_{P-C} + ^4J_{P-C}|$ .
- [27]  $N = |^1J_{P-C} + ^3J_{P-C}|$ . For details on *N* see: J. P. Fackler, Jr., J. A. Fetchin, J. Mayhew, W. C. Seidel, T. J. Swift, M. Weeks, *J. Am. Chem. Soc.* **1969**, *91*, 1941–1947.
- [28] G. M. Sheldrick, SHELXS/L-97, Programs for Crystal Structure Determination, University of Göttingen, Göttingen (Germany) **1997**. See also: G. M. Sheldrick, *Acta Crystallogr.* **1990**, *A46*, 467–473; *ibid.* **2008**, *A64*, 112–122.