

# [SbCl<sub>2</sub>( $\eta^2$ -1,4-dt)( $\mu_2$ -1,4-dt)][AlCl<sub>4</sub>]: Ionic Liquid-based Synthesis of a Stibonium Cation (dt = dithiane)

Dominic Freudenmann and Claus Feldmann

Institut für Anorganische Chemie, Karlsruhe Institute of Technology (KIT), Engesserstraße 15,  
D-76131 Karlsruhe, Germany

Reprint requests to Prof. Dr. C. Feldmann. Tel.: ++49-721-60842855.

E-mail: [claus.feldmann@kit.edu](mailto:claus.feldmann@kit.edu)

*Z. Naturforsch.* **2013**, *68b*, 1090 – 1094 / DOI: 10.5560/ZNB.2013-3133

Received May 8, 2013

The reaction of antimony(III) chloride with 1,4-dithiane (1,4-dt) in the ionic liquid [BMIM]Cl ([BMIM]Cl: 1-butyl-4-methylimidazolium chloride) with an excess of the Lewis acid AlCl<sub>3</sub> results in the formation of a ligand-stabilized stibonium cation [SbCl<sub>2</sub>]<sup>+</sup>. 1,4-Dithiane simultaneously serves as a chelating  $\eta^2$ -ligand, exhibiting a boat-like conformation, as well as a bridging  $\mu_2$ -ligand, exhibiting a chair-like conformation, with Sb<sup>3+</sup> as the coordinating cation. Due to the bridging 1,4-dithiane ligands, a chain-like structural building unit  $^1_{\infty}[\text{SbCl}_2(\eta^2\text{-1,4-dt})_{1/1}(\mu_2\text{-1,4-dt})_{2/2}]^+$  is formed.

**Key words:** Stibonium Cation, Ionic Liquid, Crystal Structure, 1,4-Dithiane

## Introduction

Electron-rich pnictogen centers are known to react as Lewis acceptors although a lone pair is present at the main-group element [1, 2]. The resulting stibonium- or stibinidenium cations in the case of antimony, as well as comparable species of other pnictogens, are typically obtained as ligand-stabilized salts and lead to a wide range of different coordination compounds of the main-group elements. Stibonium cations most often form coordination compounds with group 15 elements, *i. e.*, N-, P-, As- or Bi-containing ligands [3–5]. In contrast, stibonium cations coordinated by group 16 elements are rare to date.

As a part of our studies concerning the benefit of ionic liquids (ILs) in inorganic synthesis, we studied their use in view of a coordination of chalcogen-containing ligands to pnictogen centers [6]. Lewis-acidic ionic liquids are known as suitable reaction media for the synthesis of inorganic compounds exhibiting novel structural building units and bonding situations [7]. This also includes cases of element-element bonding and low oxidation states of the main-group elements. Based on this strategy, the title compound [SbCl<sub>2</sub>( $\eta^2$ -1,4-dt)( $\mu_2$ -1,4-dt)][AlCl<sub>4</sub>] was obtained that is constituted of a [SbCl<sub>2</sub>]<sup>+</sup> stibe-

nium cation. This stibonium cation is coordinated and interlinked by 1,4-dithiane (1,4-dt) as a ligand. Here, [SbCl<sub>2</sub>( $\eta^2$ -1,4-dt)( $\mu_2$ -1,4-dt)][AlCl<sub>4</sub>] represents the first stibonium cation that is coordinated by group 16 elements. Surprisingly, 1,4-dt acts as a bidentate bridging ligand as well as a bidentate chelating ligand. Noteworthy, only three compounds of extended network structures with 1,4-dithiane as a linker have yet been reported. These examples comprise silver(I)- and indium(III)-based coordination compounds [8–10]. A simultaneous coordination of 1,4-dt as a bridging and chelating ligand is shown with the title compound for the first time.

## Results and Discussion

[SbCl<sub>2</sub>( $\eta^2$ -1,4-dt)( $\mu_2$ -1,4-dt)][AlCl<sub>4</sub>] was prepared by reaction of antimony(III) chloride and 1,4-dithiane in the Lewis-acidic ionic liquid [BMIM][AlCl<sub>4</sub>]. The synthesis resulted in colorless crystals of the title compound. X-Ray structure analysis based on single crystals revealed [SbCl<sub>2</sub>( $\eta^2$ -1,4-dt)( $\mu_2$ -1,4-dt)][AlCl<sub>4</sub>] to crystallize monoclinically in space group *P2<sub>1</sub>/c* with *Z* = 4 (Table 1). In addition to single-crystal structure analysis, the thermal behavior of the title compound

was investigated by differential thermal analysis and thermogravimetry (DTA-TG).

According to the crystal structure analysis,  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})][\text{AlCl}_4]$  is constituted of  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})]^+$  cations and  $[\text{AlCl}_4]^-$  anions (Fig. 1). Antimony is coordinated by two chlorine atoms forming a stibonium cation  $[\text{SbCl}_2]^+$  (Fig. 2). Herein, the Sb–Cl distances are almost equal in length (237.1 and 239.6 pm) which is in agreement with values reported for comparable stibonium compounds, including the phosphane-stabilized compound  $[\text{SbCl}_2(\text{PPh}_2)_2][\text{AlCl}_4]$  (Sb–Cl: 239.0 pm) [2]. The Cl–Sb–Cl angle in the title compound is near to a rectangular arrangement ( $91.7^\circ$ ) (Table 2), whereas comparable compounds show a somewhat wider angle (*e. g.*,  $\angle(\text{Cl-Sb-Cl})$  in  $[\text{SbCl}_2(\text{PPh}_2)_2]^+$  with  $96.5^\circ$  [2]). In addition to the direct Sb–Cl bonding, long-ranging Sb–Cl<sub>Al</sub> contacts (up to 380 pm) to neighboring  $[\text{AlCl}_4]^-$  anions are observed (Fig. 3). These Sb–Cl<sub>Al</sub> contacts are below the sum of the van der Waals radii of antimony and chlorine ( $d_{\text{Sb+Cl}} = 400$  pm) [11]. The  $[\text{AlCl}_4]^-$  counterions show a mean Al–Cl distance of 211 pm and a mean Cl–Al–Cl angle of  $110^\circ$ , indicating the expected distorted tetrahedral shape (Fig. 3). These data are well in accordance with those of known tetrachloridoaluminates(III) [12]. Two chlorine atoms of the  $[\text{AlCl}_4]^-$  anion (Cl5, Cl6) show somewhat increased parameters of thermal motion (Fig. 3). This finding can be explained by two sets of triple hydrogen




Fig. 1. Unit cell of  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})][\text{AlCl}_4]$ .




Fig. 2. Zig-zag-shaped arrangement of the  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})_{1/1}(\mu_2\text{-1,4-dt})_{2/2}]^+$  cation (displacement ellipsoids with 50% probability of finding).

bonding of these two chlorine atoms with the chelating 1,4-dt ligand.

The  $[\text{SbCl}_2]^+$  stibonium cations are interlinked *via* three molecules of 1,4-dithiane (Fig. 2). Together with the chlorine atoms, thus a pseudo-octahedral coordination of the central antimony atom is observed (Fig. 3). Although ligand-stabilized stibonium cations are known, such species most often exhibit phosphorus- or nitrogen-containing ligands [13–15]. As an additional feature, the 1,4-dithiane ligand in  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})]^+$  serves as both, a bidentate bridging ligand as well as a bidentate chelating ligand (Figs. 2, 3). While the chelating 1,4-dt exhibits a boat-type conformation, the bridging 1,4-dt shows a chair-type conformation. Noteworthy, two orientations of the bridging 1,4-dt are present and cause a certain displacement of the sulfur atoms (Fig. 3). This disorder was tackled by split-atom positions that were refined with non-fixed site occupation factors, resulting in an occupation of 83% (S2A) and 17% (S2B). 1,4-dt has been frequently described to act as a bidentate, bridging ligand (*e. g.*, in  $[\text{RuCl}_2(\text{dpb})(1,4\text{-dt})]$  and  $\{[\text{Ag}_2(1,4\text{-dt})_3][\text{CF}_3\text{SO}_3]_2\}_\infty$ ) [10, 16]) and typically exhibits a chair-like conformation between two metal centers similar to the situation in the title compound. Compounds with 1,4-dt as a bidentate, chelating ligand are known for altogether five compounds [16–20]. A simultaneous combination of both types of coordination – bridging and chelating – to the best of our knowledge has not yet been observed for 1,4-dt as




Fig. 3. Structure and distances of the  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})_{1/1}(\mu_2\text{-1,4-dt})_{2/2}]^+$  cation (with displacement of the 1,4-dt ligand) and of the  $[\text{AlCl}_4]^-$  anion (all distances in pm; displacement ellipsoids with 50% probability of finding).

a ligand. This finding can be ascribed to the absence of alternative ligands for coordination as well as to the weakly coordinating properties of the ionic liquid.

The Sb–S distances confirm a coordinative bonding between  $[\text{SbCl}_2]^+$  and 1,4-dt. Thus, distances in a range of 269.5 to 323.2 pm are observed (Table 2, Fig. 3) that are significantly elongated in comparison to those in binary compounds such as  $\text{Sb}_2\text{S}_3$  (Sb–S: 249 pm) or antimony thiolates (Sb–S: 241 pm) [21, 22]. Considering the coordinative Sb–S interactions, a chain-like arrangement of the cationic building unit  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})_{1/1}(\mu_2\text{-1,4-dt})_{2/2}]^+$  becomes obvious (Fig. 2). Within these chains, the bridging 1,4-dt molecules are linking *via* their S atoms which are located at the equatorial positions of  $\text{Sb}^{3+}$ . This co-




Fig. 4. Thermal decomposition of  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})][\text{AlCl}_4]$  (total sample weight 20.1 mg).

Table 1. Crystallographic data of  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})][\text{AlCl}_4]$ .

Empirical formula	$\text{Sb}_4\text{Cl}_6\text{AlCl}_8\text{H}_{16}$
Crystal system	monoclinic
Space group	$P2_1/c$
Lattice parameters	$a = 14.879(3) \text{ \AA}$ $b = 10.825(2) \text{ \AA}$ $c = 13.509(3) \text{ \AA}$ $\beta = 109.17(3)^\circ$ $V = 2055.2(7) \text{ \AA}^3$
Formula units per cell	$Z = 4$
Calculated density	$\rho = 1.94 \text{ g cm}^{-3}$
Measurement device	image plate diffractometer IPDS II (Stoe, Darmstadt) $\lambda(\text{MoK}\alpha) = 0.71073 \text{ \AA}$ ; $T = 200 \text{ K}$
Measurement limits	$2.9^\circ \leq 2\theta \leq 58.6^\circ$ $0 \leq h \leq 20, -14 \leq k \leq 14, -18 \leq l \leq 17$ $\mu = 2.6 \text{ mm}^{-1}$
Linear absorption coefficient	
Crystal size	$0.17 \times 0.12 \times 0.04 \text{ mm}^3$
Absorption correction	numerical ( $T_{\text{min}}: 0.49; T_{\text{max}}: 0.80$ )
Number of reflections	10034 measured; thereof 5191 independent
Data averaging	$R_{\text{int}} = 0.023$
Structure refinement	full matrix least-squares on $F_0^2$ , anisotropic displacement parameters for all non-hydrogen atoms
Number of parameters	192
Figures of merit	$R1 = 0.055$ $R1(3818 F_0^2 > 2 \sigma F_0^2) = 0.041$ $wR2 = 0.120$ (all data) Goof = 0.98
Residual electron density	$-0.91 / +0.82 \text{ e \AA}^{-3}$

ordination results in a zig-zag-shaped chain. Some C–H groups of the 1,4-dt ligand are involved in various intermolecular hydrogen bonding interactions to chlorine atoms of neighboring  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-$

Table 2. Selected bond lengths and angles of the  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})]^+$  cation.

Atoms	Distance (pm)	Atoms	Angle (deg)
Sb1–Cl1	237.1(1)	Cl1–Sb1–Cl2	91.7(1)
Sb1–Cl2	239.6(1)	Cl1–Sb1–S1	84.8(1)
Sb1–S1	302.3(1)	S1–Sb1–S2	108.5(1)
Sb1–S2	323.2(1)	S4–Sb1–Cl2	82.5(1)
Sb1–S3	300.2(1)	S3–Sb1–Cl1	79.3(1)
Sb1–S4	269.5(1)	C1–S1–C2	100.5(1)

1,4-dt)]<sup>+</sup> cations and  $[\text{AlCl}_4]^-$  anions. The C–H···Cl distances range between typical values of 267 to 333 pm [23].

Finally, the thermal behavior of  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})][\text{AlCl}_4]$  was analyzed by performing thermogravimetry (Fig. 4). Upon heating to 600 °C, the title compound exhibits a significant weight loss of about 76% between 120 and 300 °C. This can be attributed to an evaporation of  $\text{SbCl}_3$  and 1,4-dithiane (calcd. 77.8%). When increasing the temperature to more than 380 °C, a second decomposition step (weight loss: 23%) occurred that can be ascribed to an evaporation of  $\text{Al}_2\text{Cl}_6$  (calcd. 22.2%). These findings match well with the respective boiling points of  $\text{SbCl}_3$  ( $T_{\text{bp}}(\text{SbCl}_3)$ : 223 °C) and the ligand ( $T_{\text{bp}}(1,4\text{-dt})$ : 200 °C). Evaporation of the  $\text{AlCl}_3$  is observed at rather high temperatures, which can be ascribed to the interaction of this strong Lewis acid with  $\text{Al}_2\text{O}_3$  as the crucible material.

## Conclusions

$[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})][\text{AlCl}_4]$  consists of  $[\text{SbCl}_2]^+$  stibonium cations that are interlinked to zig-zag chains *via* 1,4-dithiane as a ligand.  $[\text{AlCl}_4]^-$  is positioned between the  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})]_{1/1}(\mu_2\text{-1,4-dt})_{2/2}]^+$  chains as a weakly coordinating anion. In general, stibonium cations that are stabilized by chalcogen-containing ligands are rare to date. The coordinative situation of the 1,4-dt ligand is unusual since it simultaneously serves as a bidentate chelating ligand as well as a bidentate bridging ligand. Both types of coordination are known but, till now, not realized in a single compound. Notably, chelating 1,4-dt exhibits a boat-like conformation, whereas bridging 1,4-dt shows a chair-like conformation. The formation of  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})][\text{AlCl}_4]$  as a coordination compound is based on the ionic-liquid system

and its weakly coordinating properties. Perspectively, this strategy of reaction could be used to realize many more coordination compounds of electron-rich pnictogen centers with unusual bonding and coordination situation.

## Experimental Section

### Materials and synthesis

The experiments were carried out under standard Schlenk and argon glove box techniques. Reactions were performed in argon-filled and sealed glass ampoules that were dried at 300 °C before use. The commercially available starting materials antimony(III) chloride (99.99%, Sigma-Aldrich), aluminum(III) chloride (99.99%, Sigma-Aldrich) and 1,4-dithiane (97%, ABCR) were dried in vacuum for 24 h. To remove all traces of water, the commercially available ionic liquid  $[\text{BMIM}]\text{Cl}$  (Iolitec) was dried at 120 °C for two days in vacuum prior to use.

To synthesize  $[\text{SbCl}_2(\eta^2\text{-1,4-dt})(\mu_2\text{-1,4-dt})][\text{AlCl}_4]$ , 100 mg (0.44 mmol) of antimony(III) chloride, 105.4 mg (0.876 mmol) of 1,4-dithiane and 233.8 mg (1.75 mmol) of aluminum(III) chloride were reacted in the ionic liquid  $[\text{BMIM}]\text{Cl}$  at 75 °C for 5 days. Colorless crystals were obtained after cooling to room temperature with a rate of 1 K/h. These crystals were separated from the ionic liquid by filtration through a glass filter, followed by washing twice with 2 mL of dried diethyl ether. Finally, the title compound was dried in a vacuum overnight and obtained with a yield of about 70%.

### Characterization and analytical tools

For single-crystal structure analysis suitable crystals were chosen and fixed in glass capillaries with inert oil (KEL-F). Complete intensity data sets were collected with an IPDS II diffractometer (Stoe, Darmstadt) using  $\text{MoK}\alpha$  radiation ( $\lambda_\alpha$ : 71.073 pm, graphite monochromator). Data reduction was performed using the program package X-Area [24]. The structure was solved by Direct Methods using SHELXS-97 [25]. Anisotropic refinement on  $2\sigma F^2$  was carried out for all non-hydrogen atoms by full-matrix least-squares methods using the program SHELXL-97 [26]. Disorder of the bridging ligand's sulfur atoms was observed and tackled by split atom positions with 83 and 17% occupation (Fig. 3). The positions of the H atoms were modelled based on idealized C–H bonds using a riding model. A numerical absorption correction was applied using the program X-RED [27]. Crystallographic data and structure refinement details are summarized in Table 1. Selected bond lengths and angles of the title compound are listed in Table 2. Crystallographic illustrations were created with DIAMOND [28].

CCDC 937474 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

Differential thermal analysis/thermogravimetry (DTA/TG) was performed with a Netzsch STA 409C instrument applying  $\alpha$ -Al<sub>2</sub>O<sub>3</sub> as a crucible material and reference. The samples were heated under N<sub>2</sub> flow to 600 °C with a rate of 10 K min<sup>-1</sup>. The resulting data were baseline-corrected

by subtracting the results of a measurement with an empty crucible.

#### Acknowledgement

The authors are grateful to the Center for Functional Nanostructures (CFN) of the Deutsche Forschungsgemeinschaft (DFG) at the Karlsruhe Institute of Technology (KIT) for financial support.

- [1] N. Burford, P. J. Ragona, *J. Chem. Soc., Dalton Trans.* **2002**, 4307–4315.
- [2] S. S. Chitnis, B. Peters, E. Conrad, N. Burford, R. McDonald, M. J. Ferguson, *Chem. Commun.* **2011**, 47, 12331–12333.
- [3] N. L. Kilah, S. Petrie, R. Stranger, J. W. Wielandt, A. C. Willis, S. B. Wild, *Organometallics* **2007**, 26, 6106–6113.
- [4] E. Conrad, N. Burford, R. McDonald, M. J. Ferguson, *J. Am. Chem. Soc.* **2009**, 131, 5066–5067.
- [5] E. Conrad, N. Burford, R. McDonald, M. J. Ferguson, *Chem. Commun.* **2010**, 46, 4598–4600.
- [6] D. Freudenmann, S. Wolf, M. Wolff, C. Feldmann, *Angew. Chem. Int. Ed.* **2011**, 50, 11050–11069.
- [7] E. Ahmed, A. Isaeva, A. Fiedler, M. Haft, M. Ruck, *Chem. Eur. J.* **2011**, 17, 6847–6852.
- [8] J. J. H. Edema, F. van Bolhuis, A. Meetsma, R. M. Kellogg, H. Kooijman, A. L. Spek, *Inorg. Chim. Acta* **1993**, 207, 263–265.
- [9] J. Blank, H. D. Hausen, W. Schwartz, J. Weidlein, *J. Organomet. Chem.* **1993**, 443, 145–151.
- [10] N. R. Brooks, A. J. Blake, N. R. Champness, J. W. Cunningham, P. Hubberstey, S. J. Teat, C. Wilson, M. Schröder, *J. Chem. Soc., Dalton Trans.* **2001**, 2530–2538.
- [11] A. F. Holleman, N. Wiberg, *Lehrbuch der Anorganischen Chemie*, de Gruyter, Berlin **2007**.
- [12] N. C. Baenziger, A. D. Nelson, *J. Am. Chem. Soc.* **1968**, 90, 6602–6607.
- [13] A. H. Spinney, I. Korobkov, D. S. Richeson, *Chem. Commun.* **2007**, 16, 1647–1649.
- [14] C. Hering, M. Lehmann, A. Schulz, A. Villinger, *Inorg. Chem.* **2012**, 51, 8212–8224.
- [15] R. J. Wiacek, C. L. Macdonald, J. N. Jones, J. M. Pietryga, A. H. Cowley, *Chem. Commun.* **2003**, 3, 430–431.
- [16] M. O. Santiago, J. R. Sousa, I. C. N. Diogenes, L. G. F. Lopes, E. Meyer, E. E. Castellano, J. Ellena, A. A. Batista, I. S. Moreira, *Polyhedron* **2006**, 25, 1543–1548.
- [17] R. D. Adams, L. Chen, J. H. Yamamoto, *Inorg. Chim. Acta* **1995**, 229, 47–54.
- [18] M. H. Johansson, H. P. Engelbrecht, *Acta Crystallogr.* **2001**, E57, 114–116.
- [19] M. Green, M. Draganjac, Y. Jiang, P. M. Nave, A. W. Cordes, C. D. Bryan, J. K. Dixon, S. L. Folkert, C.-H. Yu, *J. Chem. Crystallogr.* **2003**, 33, 473–479.
- [20] U. Frey, S. Elmroth, B. Moullet, L. I. Elding, A. E. Merbach, *Inorg. Chem.* **1991**, 30, 5033–5037.
- [21] P. Arun, A. G. Vedeshwar, *J. Mater. Sci.* **1996**, 31, 6507–6510.
- [22] C. A. Allen, V. M. Cangelosi, L. N. Zakharov, D. W. Johnson, *Cryst. Growth Des.* **2009**, 9, 3011–3013.
- [23] H. Zhang, J. Jiang, N. Yang, H. Dang, B. K. Teo, *J. Cluster Sci.* **1998**, 9, 547–554.
- [24] X-AREA, X-RED (version 1.26), Area Detector Software System, Stoe & Cie GmbH, Darmstadt (Germany) **2004**.
- [25] G. M. Sheldrick, *Acta Crystallogr.* **1990**, A46, 467–473.
- [26] G. M. Sheldrick, *Acta Crystallogr.* **2008**, A64, 112–122.
- [27] X-RED, X-SHAPE (version 1.26), Crystal Optimization for Numerical Absorption Correction, Stoe & Cie GmbH, Darmstadt (Germany) **2004**.
- [28] K. Brandenburg, DIAMOND (version 3.2i), Crystal and Molecular Structure Visualization, Crystal Impact – K. Brandenburg & H. Putz GbR, Bonn (Germany) **2012**. See also: <http://www.crystalimpact.com/diamond/>.